

Environmental Protection (Water) Policy 2009

Proserpine River, Whitsunday Island and O'Connell River Basins Environmental Values and Water Quality Objectives

Basins:

122, including all waters of the Proserpine River Basin
and adjacent coastal waters

123, including all waters of the Whitsunday Island Basin
and adjacent coastal waters

124, including all waters of the O'Connell River Basin and
adjacent coastal waters



Prepared by: Environmental Policy and Planning Division, Department of Environment and Heritage Protection

© State of Queensland, 2013.

The Queensland Government supports and encourages the dissemination and exchange of its information. The copyright in this publication is licensed under a Creative Commons Attribution 3.0 Australia (CC BY) licence.



Under this licence you are free, without having to seek our permission, to use this publication in accordance with the licence terms.

You must keep intact the copyright notice and attribute the State of Queensland as the source of the publication.


For more information on this licence, visit <http://creativecommons.org/licenses/by/3.0/au/deed.en>

If you need to access this document in a language other than English, please call the Translating and Interpreting Service (TIS National) on 131 450 and ask them to telephone Library Services on +61 7 3170 5470.

This publication can be made available in an alternative format (e.g. large print or audiotape) on request for people with vision impairment; phone +61 7 3170 5470 or email library@ehp.qld.gov.au.

August 2013

Main parts of this document and what they contain



Contents

Main parts of this document and what they contain	i
1 Introduction	1
1.1 Waters to which this document applies.....	1
1.2 Guidance on using this document.....	2
1.3 Information about mapped areas and boundaries	4
1.4 Water types and basis for boundaries	4
1.5 Matters for amendment.....	5
2 Environmental values	6
2.1 Environmental values.....	6
2.2 Management goals to support environmental values	6
3 Water quality objectives to protect environmental values	12
3.1 Water quality objectives to protect aquatic ecosystems	12
3.2 Water quality objectives for human use environmental values.....	51
4 Ways to improve water quality.....	65
5 Dictionary.....	66

List of tables

Table 1. Environmental values for Proserpine River, Whitsunday Island and O'Connell River Basin waters	7
Table 2. Water quality objectives to protect aquatic ecosystem environmental value under ambient (baseflow) and event (high flow) conditions.....	14
Table 3. Water quality objectives to protect human use environmental values.	51
Table 4. Drinking water EV: Priority water quality objectives for drinking water supply in the vicinity of off-takes, including groundwater, before treatment.....	54
Table 5. Aquaculture EV: Water quality objectives for tropical aquaculture.....	56
Table 6. Aquaculture EV: Water quality objectives for optimal growth of particular species in fresh water.....	57
Table 7. Aquaculture EV: Water quality objectives for optimal growth of particular marine species.	58
Table 8. Irrigation EV: Water quality objectives for thermotolerant (faecal) coliforms in irrigation waters used for food and non-food crops.	59
Table 9. Irrigation EV: Water quality objectives for heavy metals and metalloids in agricultural irrigation water— long-term trigger value (LTV), short-term trigger value (STV) and soil cumulative contamination loading limit (CCL).	60
Table 10. Stock watering EV: Water quality objectives for tolerances of livestock to total dissolved solids (salinity) in drinking water.	61
Table 11. Stock watering EV: Water quality objectives (low risk trigger values) for heavy metals and metalloids in livestock drinking water.	62
Table 12. Recreational waters: Alert levels and corresponding actions for management of cyanobacteria.....	63
Table 13. Suite of environmental values that can be chosen for protection.....	67

1 Introduction

This document is made pursuant to the provisions of the Environmental Protection (Water) Policy 2009 (EPP (Water)), which is subordinate legislation under the *Environmental Protection Act 1994*. The EPP (Water) provides a framework for:

- identifying environmental values (EVs) for Queensland waters, and deciding the water quality objectives (WQOs) to protect or enhance those EVs. (WQOs are long term goals for receiving waters, not individual point source emission objectives.)
- including the identified EVs and WQOs under Schedule 1 of the EPP (Water).
- This document contains EVs and WQOs for waters in the Proserpine River, Whitsunday Island and O'Connell River basins and adjacent coastal waters, and is listed under schedule 1 of the EPP (Water).

1.1 Waters to which this document applies

This document applies to fresh and estuarine surface waters and groundwaters draining the basins of the Proserpine River, Whitsunday Island, and O'Connell River (basins 122, 123 and 124¹), and coastal waters as indicated in the accompanying plans (WQ1221—surface waters, WQ1222—coastal waters)².

Waters covered by this document include:

- Proserpine River Basin (excluding those waters within Abbot Point port limits), including fresh and estuarine waters of the Proserpine River, Eden Lassie Creek, Gregory River, Thompson, Myrtle and Repulse creeks, and Lethe Brook
- O'Connell River Basin, including fresh and estuarine waters of the O'Connell and Andromache rivers, Waterhole, Blackrock, St Helens, Murray, Constant, and Reliance creeks. (A small southern section of the O'Connell River Basin near Mackay has been included with the Pioneer River Basin EVs document.)
- Whitsunday Islands, including fresh and estuarine waters
- wetlands, lakes and reservoirs
- groundwaters
- waters on islands
- enclosed coastal and open coastal waters, excluding Edgecumbe Bay and other waters within Abbot Point port limits.

The geographical extent of waters addressed by this document is shown in plans WQ1221 and WQ1222, and is broadly:

- north to the boundary of the Proserpine River Basin with the Abbot Point port limits
- west to the boundary with the Don and Burdekin river basins
- south to and including the Reliance Creek subcatchment
- east and north-east to the limit of Queensland Coastal Waters.

¹ Australia's River Basins 1997—Product User Guide. Published by Geoscience Australia. Canberra, ACT (3rd edition, 2004).

² This document and the accompanying plans are available from the department's website at www.ehp.qld.gov.au. The boundaries in the accompanying plans WQ1221 and WQ1222 are indicative only. EVs, water types and aquatic ecosystem management intent (level of protection) depicted in the accompanying plans are stored in electronic form as part of the Central Queensland Environmental Values Schedule 1 Geodatabase August 2013, and held at the department's offices at 400 George Street Brisbane. Geodatabase regions are based on the regions established in the Queensland Water Quality Guidelines. Spatial (GIS) datasets can be downloaded free of charge from the Queensland Government Information Service (QGIS) at <http://dds.information.qld.gov.au/dds>. For further information, email the department at epa.ev@ehp.qld.gov.au.

1.2 Guidance on using this document

1.2.1 Key terms (refer to dictionary for additional terms)

ADWG means the Australian Drinking Water Guidelines (2011), prepared by the National Health and Medical Research Council (NHMRC) in collaboration with the Natural Resource Management Ministerial Council (NRMMC)³.

AWQG means the Australian and New Zealand Guidelines for Fresh and Marine Water Quality (October 2000), prepared by the Australian and New Zealand Environment and Conservation Council (ANZECC) and the Agriculture and Resource Management Council of Australia and New Zealand (ARMCANZ)⁴.

Environmental values (EVs) for water means the EVs specified in Table 1 of this document for the corresponding water.

EVs for water are the qualities of water that make it suitable for supporting aquatic ecosystems and human water uses. These EVs need to be protected from the effects of habitat alteration, waste releases, contaminated runoff and changed flows to ensure healthy aquatic ecosystems and waterways that are safe for community use. Particular waters may have different EVs. The range of EVs and the waters they can potentially apply to are listed below, and further details are provided in the dictionary (refer section 5).

List of EVs and applicable waters

Environmental value (EV)	Potentially applicable to:	
	Tidal waters	Fresh (non-tidal) waters
<p>Protection of aquatic ecosystems (aquatic ecosystem EV)</p> <p>Protection or enhancement of aquatic ecosystem values, under four possible levels of ecosystem conditions:</p> <ul style="list-style-type: none"> • high ecological value (effectively unmodified) waters • slightly disturbed waters • moderately disturbed waters • highly disturbed waters. <p>(Suitability for seagrass and wildlife habitat have also been specifically identified for some Queensland waters as a component of this EV).</p>	✓	✓
<p>EVs other than aquatic ecosystem EV (called human use EVs)</p> <p>Suitability for drinking water supplies</p> <p>Suitability for primary contact recreation (e.g. swimming)</p> <p>Suitability for secondary contact recreation (e.g. boating)</p> <p>Suitability for visual (no contact) recreation</p> <p>Suitability for human consumers of wild or stocked fish, shellfish or crustaceans (suitability for oystering has also been specifically identified for some Queensland waters)</p> <p>Protection of cultural and spiritual values, including traditional owner values of water</p> <p>Suitability for industrial use (including mining, minerals refining/processing)</p> <p>Suitability for aquaculture (e.g. red claw, barramundi)</p> <p>Suitability for crop irrigation</p> <p>Suitability for stock watering</p> <p>Suitability for farm supply/use</p>	✓ ✓ ✓ ✓ ✓ ✓ ✓ ✓ ✓ ✓ ✓	✓ ✓ ✓ ✓ ✓ ✓ ✓ ✓ ✓ ✓ ✓

³ The ADWG are available on the National Health and Medical Research Council website at www.nhmrc.gov.au.

⁴ The AWQG are available on the Australian Government's National Water Quality Management Strategy website.

Level of protection for a water (aquatic ecosystem EV) means the level of aquatic ecosystem condition specified in Table 2 of this document that the corresponding WQOs for that water are intended to achieve (refer to management intent definition below for further information).

Management goal means the goals (if any) stated in section 2 of this document to support the EVs for waters identified in Table 1.

Management intent (level of protection) for a water (aquatic ecosystem EV) means the level of aquatic ecosystem condition specified in Table 2 of this document that the corresponding WQOs for that water are intended to achieve. For example, the intent for high ecological value waters is that their effectively unmodified condition is maintained.

QWQG means the Queensland Water Quality Guidelines.⁵

Water quality guidelines (defined in the EPP (Water)) are numerical concentration levels or statements for indicators that protect a stated environmental value. Under the EVs setting process contained in the EPP (Water), water quality guidelines are used as an input to the development of WQOs.

Water quality indicator (for an EV) means a property that is able to be measured or decided in a quantitative way. Examples of water quality indicators include physical indicators (e.g. temperature), chemical indicators (e.g. nitrogen, phosphorus, metals), and biological indicators (e.g. macroinvertebrates, seagrass, fish).

Water quality objectives (WQOs) means the WQOs specified in tables 2–12 of this document to support the EVs for waters identified in Table 1.

WQOs are long-term goals for water quality management. They are numerical concentration levels or narrative statements of indicators established for receiving waters to support and protect the designated EVs for those waters. Water quality objectives are not individual point source emission objectives, but the receiving water quality objectives.

They are based on scientific criteria or water quality guidelines but may be modified by other inputs (e.g. social, cultural, economic).

Examples of WQOs include:

- total phosphorus concentration less than 20 micrograms per litre (µg/L)
- chlorophyll a concentration less than 1 µg/L
- dissolved oxygen between 95 per cent and 105 per cent saturation
- family richness of macroinvertebrates greater than 12 families
- exotic individuals of fish less than five per cent.

Water type means groupings of waters with similar characteristics, as shown in the accompanying plans. The water types covered by this document are based on mapping and definitional rules for water types established in the QWQG and, where available, other site-specific studies and documents. Water types can include fresh waters (lowland, upland, lakes/reservoirs), wetlands and groundwaters, estuarine waters (lower, middle and upper estuaries), tidal canals, constructed estuaries, marinas and boat harbours, and coastal marine waters (open coastal, enclosed coastal). WQOs applying to different water types are outlined in this document. More detail on water types is provided in section 1.4.

Refer to dictionary for additional terms.

1.2.2 Main components of this document

The main components of this document are:

- Plan WQ1221—showing the spatial extent and boundaries of surface water types covered by this document
- Plan WQ1222—showing the spatial extent and boundaries of coastal water types covered by this document
- Section 1—introduction and guidance on how to use the document
- Section 2 (Table 1)—EVs applying to waters covered by this document
- Section 3 (Tables 2–12)—WQOs applying to different EVs:
 - Table 2 provides WQOs to protect the aquatic ecosystem EV, and closely links to the water types shown on

⁵ The QWQG are available on the department's website.

plan WQ1221, and plan WQ1222 (coastal waters)

- o tables 3 to 12 provide WQOs to protect human use EVs
- Section 4—ways to improve water quality: containing a list of relevant documents, provided for information purposes only
- Section 5—a dictionary of other terms relevant to EVs and WQOs.

1.2.3 Use of this document

Section 2 (Table 1) lists the identified EVs for protection for particular waters. The aquatic ecosystem EV is a default applying to all Queensland waters. Reference to section 3 (Table 2) provides the corresponding WQOs to protect the aquatic ecosystem EV. Where relevant, different WQOs are specified to protect the aquatic ecosystem EV in different water types (refer to the tables and the accompanying plans). For the human use EVs specified in Table 1, tables 3 onwards provide the corresponding WQOs to support these EVs.

Where reference to Table 1 indicates more than one EV applies to a given water, the adoption of the most stringent WQO for the identified EVs applies to each water quality indicator in order to protect all identified EVs. Further detail on selection of most stringent WQOs is provided in section 3.

This document also refers to a number of guidelines, codes and other reference sources on water quality. In particular, the QWQG prepared by the department provide a technical basis for the WQOs contained in this document. The QWQG also provide more detailed information on water types, water quality indicators, derivation of local water quality guidelines, application during flood events, monitoring, and predicting and assessing compliance.

1.3 Information about mapped areas and boundaries

The boundaries in the accompanying plans WQ1221 and WQ1222 are indicative only. EVs, water types and aquatic ecosystem management intent (level of protection) depicted in the accompanying plans are stored in electronic form as part of the Central Queensland Environmental Values Schedule 1 Geodatabase August 2013, and held at the department's offices at 400 George Street Brisbane. Geodatabase regions are based on the regions established in the QWQG. Spatial (GIS) datasets can be downloaded free of charge from the Queensland Government Information Service (QGIS) at <http://dds.information.qld.gov.au/dds>. For further information, email the department at epa.ev@ehp.qld.gov.au.

1.4 Water types and basis for boundaries

1.4.1 Water types

Waters in this document have been classified into the following different water types (not all water types are present in all areas):

- Fresh waters— freshwater streams and rivers
- freshwater lakes/reservoirs
- groundwaters
- mid estuary—waters extending the majority of the length of estuaries with a moderate amount of water movement from either freshwater inflow or tidal exchange
- enclosed coastal/lower estuary—waters occurring at the downstream end of estuaries and including shallow coastal waters in adjacent enclosed bays
- marinas, boat harbours, tidal canals, and constructed estuaries,
- wetlands
- open coastal waters—waters extending to the seaward limits of Queensland waters.

The water types are based on local water quality studies in the Proserpine River, Whitsunday Island and O'Connell River basins (refer to the source documents listed after Table 2), the AWQG and mapping and definitional rules contained in the QWQG. Further detail on water types is contained in these sources.

Water types identified in this document are shown in Table 2 and the accompanying plans (WQ1221, WQ1222).

1.4.2 Water type boundaries

The boundaries of different water types have been mapped using a variety of attributes, including:

1. geographic coordinates
2. catchment or subcatchment boundaries
3. highest/lowest astronomical tide
4. tidal limiting structure (weirs)
5. maritime mapping conventions
6. coastline
7. surveyed terrestrial boundaries
8. altitude.

The basis of different boundaries is shown in the plan. The boundaries of water types may be confirmed or revised by site investigations. Refer to section 1.3 above.

1.5 Matters for amendment

Amendments of the following type may be made to this schedule 1 document for the purposes of replacement under section 12(2)(b) of the EPP (Water):

- changes to EVs
- changes to management goals
- changes to WQOs
- changes to management intent (level of protection) categories
- changes to waterway or water type boundaries/descriptions
- updates to information/data sources, websites and email contact details, agency/departmental names, other institutional names, references.

2 Environmental values

2.1 Environmental values

Table 1 and the accompanying plans WQ1221 and WQ1222 outline the EVs for waters in the Proserpine River, Whitsunday Island and O'Connell River basins, and adjacent coastal waters. These are based on stakeholder EVs consultations undertaken by the department and Mackay Whitsunday NRM (Reef Catchments) as part of the Mackay Whitsunday region water quality improvement plan, and additional consultation by the department in preparing this document. Consultation results from the water quality improvement plan are reported in:

- Mitchell, C & Higham, W 2008. Water quality improvement plan (WQIP): Community consultation, Mackay Whitsunday Natural Resource Management Group, Reef Catchments, Mackay.

The dictionary to this document provides further explanation of EVs (refer section 5).

2.2 Management goals to support environmental values

There are no management goals specified under this document.

Table 1. Environmental values for Proserpine River, Whitsunday Island and O'Connell River Basin waters

	Environmental values ^{1,2,3,4,5}											
	Aquatic ecosystems	Irrigation	Farm supply/use	Stock water	Aquaculture	Human consumer	Primary recreation ⁵	Secondary recreation ⁵	Visual recreation	Drinking water	Industrial use	Cultural and spiritual values
Subcatchment / waterway name (alphabetically listed—fresh water, estuarine, coastal/marine)												
Andromache River fresh waters	✓	✓	✓	✓		✓	✓		✓	✓		✓
Blackrock Creek fresh waters	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓		✓
Blackrock Creek estuarine waters	✓					✓		✓	✓			✓
Coastal / marine waters - refer to 'Whitsunday Islands' and 'Southern Cumberland Islands' rows												
Constant Creek fresh waters	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓		✓
Constant Creek estuarine waters	✓					✓		✓	✓			✓
Eden Lassie Creek fresh waters (outside Abbot Point port limits)	✓	✓	✓	✓	✓	✓			✓	✓		✓
Eden Lassie Creek estuarine waters (outside Abbot Point port limits)	✓				✓	✓		✓	✓			✓
Gregory River fresh waters (outside port limits)	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓
Gregory River estuarine waters (outside port limits)	✓				✓	✓		✓	✓			✓
Island waters - refer to 'Whitsunday Islands', 'Cumberland Islands' rows												

Proserpine River, Whitsunday Island and O'Connell River Basins Environmental Values and Water Quality Objectives

	Environmental values ^{1,2,3,4,5}											
	Aquatic ecosystems	Irrigation	Farm supply/use	Stock water	Aquaculture	Human consumer	Primary recreation ⁵	Secondary recreation ⁵	Visual recreation	Drinking water	Industrial use	Cultural and spiritual values
Subcatchment / waterway name (alphabetically listed—fresh water, estuarine, coastal/marine)												
Lethe Brook fresh waters	✓	✓	✓	✓		✓	✓		✓	✓		✓
Lethe Brook estuarine waters	✓					✓		✓	✓			✓
Murray Creek fresh waters	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓		✓
Murray Creek estuarine waters	✓					✓		✓	✓			✓
Myrtle Creek fresh waters	✓	✓	✓	✓		✓	✓		✓	✓	✓	✓
O'Connell River fresh waters	✓	✓	✓	✓	✓	✓	✓		✓	✓		✓
O'Connell River estuarine waters	✓				✓	✓		✓	✓			✓
Proserpine River fresh waters	✓	✓	✓	✓		✓	✓		✓	✓	✓	✓
Proserpine River estuarine waters	✓					✓		✓	✓			✓
Reliance Creek fresh waters	✓	✓	✓	✓		✓	✓	✓	✓	✓		✓
Reliance Creek estuarine waters	✓				✓	✓		✓	✓			✓
Repulse Creek fresh waters	✓					✓	✓		✓	✓		✓

Proserpine River, Whitsunday Island and O'Connell River Basins Environmental Values and Water Quality Objectives

	Environmental values ^{1,2,3,4,5}											
	Aquatic ecosystems	Irrigation	Farm supply/use	Stock water	Aquaculture	Human consumer	Primary recreation ⁵	Secondary recreation ⁵	Visual recreation	Drinking water	Industrial use	Cultural and spiritual values
Subcatchment / waterway name (alphabetically listed—fresh water, estuarine, coastal/marine)												
Repulse Creek estuarine waters	✓					✓		✓	✓			✓
Repulse Bay	✓					✓	✓	✓	✓			✓
Southern Cumberland Islands fresh waters	✓	✓ ⁷						✓	✓	✓ ⁷		✓
Southern Cumberland Islands coastal and marine waters	✓					✓	✓	✓	✓	✓ ⁶		✓
St Helens Creek fresh waters	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓		✓
St Helens Creek estuarine waters	✓					✓		✓	✓			✓
St Helens Bay	✓					✓	✓	✓	✓			✓
Thompson Creek fresh waters	✓	✓	✓	✓		✓	✓		✓	✓		✓
Thompson Creek estuarine waters	✓					✓		✓	✓			✓
Upper Proserpine River fresh waters	✓	✓	✓	✓		✓	✓	✓	✓	✓		✓
• Lake Proserpine (Peter Faust Dam)	✓	✓		✓		✓	✓	✓	✓	✓		✓
Waterhole Creek fresh waters	✓	✓	✓	✓		✓	✓		✓	✓	✓	✓

Proserpine River, Whitsunday Island and O'Connell River Basins Environmental Values and Water Quality Objectives

	Environmental values ^{1,2,3,4,5}											
	Aquatic ecosystems	Irrigation	Farm supply/use	Stock water	Aquaculture	Human consumer	Primary recreation ⁵	Secondary recreation ⁵	Visual recreation	Drinking water	Industrial use	Cultural and spiritual values
Subcatchment / waterway name (alphabetically listed—fresh water, estuarine, coastal/marine)												
Waterhole Creek estuarine waters	✓					✓		✓	✓			✓
Whitsunday coastal creeks fresh waters	✓	✓	✓	✓		✓	✓		✓	✓		✓
Whitsunday coastal creeks estuarine waters	✓					✓		✓	✓	✓ ⁶		✓
Whitsunday Islands fresh waters ⁷ (including Whitsunday, Lindeman groups)	✓	✓ ⁷						✓	✓	✓ ⁷		✓
Whitsunday Islands estuarine waters ⁷ (including Whitsunday, Lindeman groups)	✓					✓		✓	✓	✓ ⁶		✓
Whitsunday Islands coastal and marine waters (including Whitsunday, Lindeman groups, Circular Quay Reef) - see plan WQ1222	✓					✓	✓	✓	✓	✓ ⁶		✓
Groundwaters	✓	✓	✓	✓						✓	✓	✓

Notes:

1. Refer to the accompanying plans WQ1221 and WQ1222 for locations of EVs. For fresh water and estuarine rows, the EVs shown relate to waters within each subcatchment. For example the EVs for 'Andromache River fresh waters' apply to all riverine freshwaters in the Andromache River subcatchment.
2. ✓ means the EV is selected for protection. Blank indicates that the EV is not chosen for protection.
3. Refer to the dictionary for further explanation of EVs.
4. Refer to section 3 for WQOs applying to the EVs in this table.
5. The selection of recreational EVs for waters does not mean that these waters are free of venomous or dangerous aquatic organisms, for example venomous organisms (e.g. marine stingers including box jellyfish, irukandji jellyfish), crocodiles, and sharks. Direct contact with venomous or dangerous aquatic organisms should be avoided. Refer to council websites, www.health.qld.gov.au, www.beachsafe.org.au,

Proserpine River, Whitsunday Island and O'Connell River Basins Environmental Values and Water Quality Objectives

www.marinestingers.com.au and other information sources for further details on swimming safety and information on specific waters.

6. In some areas (particularly islands) reverse osmosis plants may be in use to treat sea water for drinking water.

7. This is a generic list of EVs for islands. EVs for specific islands may differ according to degree of development, available facilities and management intent for individual islands.

3 Water quality objectives to protect environmental values

This section provides WQOs to support and protect different EVs identified for waters within the Proserpine River, Whitsunday Island and O'Connell River basins in Table 1. WQOs are long-term goals for water quality management. They are numerical concentration levels or narrative statements of indicators established for receiving waters to support and protect the designated EVs for those waters. Water quality objectives are not individual point source emission objectives, but the receiving water quality objectives.

This section is in two main parts:

- Section 3.1 (Table 2) outlines WQOs to protect the aquatic ecosystem EV. The aquatic ecosystem EV is a default applying to all Queensland waters, and therefore the WQOs for aquatic ecosystems form the minimum WQOs for all waters. Where no human use EVs are identified, the WQOs identified for aquatic ecosystem protection remain applicable.
- Section 3.2 (tables 3 to 12) provides WQOs for EVs other than aquatic ecosystem ('human use EVs') such as recreational water use, irrigating crops, and aquaculture.

Sources used in deriving WQOs are provided after the tables.

Reference to the identified EVs in Table 1 of this document provides guidance on the EVs applying to waters within the catchment. Where reference to Table 1 indicates more than one EV applies to a given water (for example aquatic ecosystem and recreational use), the most stringent WQO for each water quality indicator applies, which will then protect all identified EVs. Refer to the two following examples on selection of most stringent WQOs. Note that these are examples only and should not be directly adopted for use.

Example 1

For freshwater streams with aquatic ecosystem and drinking water EVs, the respective turbidity WQOs are:

- aquatic ecosystem lowland freshwater stream: less than 10 nephelometric turbidity units (NTU)
- drinking water: less than 25 NTU.

In this case, the aquatic ecosystem WQO for turbidity (less than 10 NTU) is the more stringent, and its adoption therefore supports both the aquatic ecosystem and drinking water EVs.

Example 2

In the following situation there are stock watering and irrigation EVs, with differing WQOs for thermotolerant (faecal) coliforms (measured as median number of organisms per 100 millilitre (mL)):

- stock watering: less than 100 organisms per 100 mL
- raw human food crops in direct contact with irrigation water: less than 10 organisms per 100 mL
- pasture and fodder for dairy animals: less than 100 organisms per 100 mL.

The most stringent WQO for faecal coliforms in this example is that for direct irrigation of raw human food crops (less than 10 organisms per 100 mL) and its adoption would in turn provide faecal coliform WQOs that protect all the above-identified human use EVs.

3.1 Water quality objectives to protect aquatic ecosystems

This section provides physico-chemical, biological (section 3.1.1) and riparian (section 3.1.2) WQOs to support the aquatic ecosystem EV. Sources used in deriving locally relevant WQOs are provided after the tables in each of these sections.

Section 5 and Appendix D of the QWQG address procedures for the application of guidelines for aquatic ecosystem protection, and compliance assessment protocols. For the comparison of test site monitoring data against WQOs, the median water quality value (e.g. concentration) of a number (preferably five or more) independent samples at a particular monitoring ('test') site should be compared against the water quality objective of the same indicator, water type and level of aquatic ecosystem protection, as listed in Table 2 below. For WQOs based on GBRMPA data, refer to notes after tables, and protocols contained in *Water quality guidelines for the Great Barrier Reef Marine Park 2010*.

3.1.1 Physico–chemical and biological water quality objectives

Table 2 includes the following information:

- water area or water type (column 1) (for boundaries of specified areas, refer to the accompanying plan)
- the corresponding management intent (level of protection) for the identified waters (column 2)
- the corresponding physico-chemical and biological WQOs to achieve the management intent (level of protection) for the identified waters.

The EPP (Water) s. 14 identifies the management intent (level of protection) for different waters. The framework and how it is addressed in this document are summarised below.

- The EPP (Water) identifies some waters for which the management intent (level of protection) is to maintain or achieve an effectively unmodified waterway condition ('high ecological value'—HEV). These may include waters that are currently HEV, 'slightly disturbed' (SD), or potentially, more modified waters which can be progressively improved to achieve HEV condition.
- In this document, waters for which all physico-chemical WQOs (e.g. nutrients, toxicants) have been set corresponding to HEV management intent are identified in columns 1 and 2 of Table 2. Each of these waters is given a specific label in the table (e.g. 'HEV1234' for effectively unmodified waters, 'SD1234' for slightly disturbed waters) which links to labels and cross-hatching on the accompanying plans.
- The management intent (level of protection) for most waters other than HEV or SD is to achieve a 'moderately disturbed' (MD) condition, for which corresponding WQOs have been derived. For some indicators and water types, WQOs correspond with a 'slightly to moderately disturbed' (SMD) level of protection, based on management intent categories specified in source technical guidelines, in particular the ANZECC (2000) Australian water quality guidelines. For ease of interpretation, this document and accompanying mapping include these within the MD level of protection. For some MD waters a higher level of protection may be provided for toxicants (e.g. pesticides).
- The management intent (level of protection) for 'highly disturbed' waters is that they be progressively improved. Some highly disturbed waters may require a long timeframe to return to a moderately disturbed condition level. In some circumstances, interim WQOs that reflect a highly disturbed (HD) condition level (which is an improvement on current condition) may be determined for such waters. Any such locations and their corresponding management intent (level of protection) are also identified in the table and accompanying plan.
- Some objectives apply to specific areas or water types as indicated in Table 2 and shown on plans WQ1221 and WQ1222, while others apply to more than one water type, as indicated in the table. The table includes ambient (baseflow) water quality objectives, and also event (high flow) objectives.

Table 2. Water quality objectives to protect aquatic ecosystem environmental value under ambient (baseflow) and event (high flow) conditions.

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
SURFACE FRESH WATERS - listed alphabetically by subcatchment (refer plan WQ1221), management intent and flow condition (ambient/event)		
Andromache River subcatchment fresh waters in area HEV2309	Aquatic ecosystem—high ecological value (HEV)	<p>Ambient (baseflow)</p> <p>Maintain existing water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas.</p> <p>The 20th, 50th and 80th percentiles of existing water quality for Andromache River subcatchment waters are:</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: 9–18–46 µg/L^{a, b, c, d} (micrograms/Litre) • particulate N: 21–39–62 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): 12–22–28 µg/L^{a, c} • particulate phosphorus: 4–9–13 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 50%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 90%–105% saturation^{a, e, f} • suspended solids: 0–1–1 mg/L^a • pH: 8.0–8.2–8.3^{a, g, h} • electrical conductivity (EC): 320–483–600 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: nd^k • fish: nd^k <p>WQOs for pesticides identified in the Mackay Whitsunday Water Quality Improvement Plan (WQIP), to protect freshwater species at the HEV level of protection^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row
Andromache River subcatchment fresh waters	Aquatic ecosystem—moderately disturbed (MD)	<p>Ambient (baseflow)</p> <p>Maintain existing water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas.</p> <p>The 20th, 50th and 80th percentiles of existing water quality for Andromache River subcatchment waters are:</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: 9–18–46 µg/L^{a, b, c, d} (micrograms/Litre) • particulate N: 21–39–62 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): 12–22–28 µg/L^{a, c} • particulate phosphorus: 4–9–13 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 50%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 90%–105% saturation^{a, e, f} • suspended solids: 0–1–1 mg/L^a • pH: 8.0–8.3^{a, g, h} • electrical conductivity (EC): 320–483–600 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: refer to 'Fresh waters - macroinvertebrates' row (listed after 'Whitsunday islands' in this table) • fish: nd^k

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
		<p>WQOs for pesticides identified in the Mackay Whitsunday WQIP^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row
<p>Andromache River subcatchment (end of system)</p>	<p>Aquatic ecosystem—moderately disturbed (MD)</p>	<p>Event (high flow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <300 µg/L^{a, b, d} • particulate N: <340 µg/L^{a, d} • filterable reactive phosphorus (FRP): <27 µg/L^a • particulate phosphorus: <70 µg/L^a • suspended solids: <200mg/L^a <p>Event WQOs for pesticides^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <0.02 µg/L • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD
<p>Blackrock Creek subcatchment fresh waters in area HEV2311</p>	<p>Aquatic ecosystem—high ecological value (HEV)</p>	<p>Ambient (baseflow)</p> <p>Maintain existing water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas.</p> <p>The 20th, 50th and 80th percentiles of existing water quality for Blackrock Creek subcatchment waters are:</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: 9–18–46 µg/L^{a, b, c, d} • particulate N: 21–39–62 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): 12–22–28 µg/L^{a, c} • particulate phosphorus: 4–9–13 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 50%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 90%–105% saturation^{a, e, f} • suspended solids: 0–1–1mg/L^a • pH: 8.0–8.2–8.3^{a, g, h} • electrical conductivity (EC): 320–430–600 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: nd^k • fish: nd^k <p>WQOs for pesticides identified in the Mackay Whitsunday WQIP, to protect freshwater species at the HEV level of protection^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
<p>Blackrock Creek subcatchment fresh waters in area SD2381 (nb: the majority of SD2381 is estuarine)</p>	<p>Aquatic ecosystem—high ecological value (HEV)</p>	<p>Ambient (baseflow) Achieve effectively unmodified water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas. Note: there is insufficient information available to establish unmodified water quality for these waters. Refer to QWQG for details on how to establish a minimum water quality data set for deriving local 20th, 50th and 80th percentiles.</p>
<p>Blackrock Creek subcatchment fresh waters</p>	<p>Aquatic ecosystem—moderately disturbed (MD)</p>	<p>Ambient (baseflow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <10 µg/L^{a, b, c, d} • particulate N: <142 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): <6 µg/L^{a, c} • particulate phosphorus: <20 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 40%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 85%–120% saturation^{a, e, f} • suspended solids: <4 mg/L^a • pH: 7.6—7.9^{a, g, h} • electrical conductivity (EC): <697 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: refer to 'Fresh waters - macroinvertebrates' row (listed after 'Whitsunday islands' in this table) • fish: nd^k <p>WQOs for pesticides identified in the Mackay Whitsunday WQIP^a:</p> <ul style="list-style-type: none"> • Ametryn: <0.02 µg/L • Atrazine: <LOD • Diuron: <0.07 µg/L • Hexazinone: <0.13 µg/L • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row
<p>Blackrock Creek subcatchment (end of system)</p>	<p>Aquatic ecosystem—moderately disturbed (MD)</p>	<p>Event (high flow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <300 µg/L^{a, b, d} • particulate N: <263 µg/L^{a, d} • filterable reactive phosphorus (FRP): <30 µg/L^a • particulate phosphorus: <70 µg/L^a • suspended solids: <33 mg/L^a <p>Event WQOs for pesticides^a:</p> <ul style="list-style-type: none"> • Ametryn: <0.06 µg/L • Atrazine: <0.55 µg/L • Diuron: <0.91 µg/L • Hexazinone: <0.37 µg/L • Tebuthiuron: <LOD

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
<p>Constant Creek subcatchment fresh waters in area HEV2314</p>	<p>Aquatic ecosystem—high ecological value (HEV)</p>	<p>Ambient (baseflow) Maintain existing water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas. The 20th, 50th and 80th percentiles of existing water quality for Constant Creek subcatchment waters are:</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: 8–11–17 µg/L^{a, b, c, d} • particulate N: 21–32–81 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): 4–5–9 µg/L^{a, c} • particulate phosphorus: 3–4–5 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 50%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 90%–105% saturation^{a, e, f} • suspended solids: 0–1–1mg/L^a • pH: 7.4–7.6–7.8^{a, g, h} • electrical conductivity (EC): 60–60–70 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: nd^k • fish: nd^k <p>WQOs for pesticides identified in the Mackay Whitsunday WQIP, to protect freshwater species at the HEV level of protection^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row
<p>Constant Creek subcatchment fresh waters in area SD2381 (nb: the majority of this SD area is estuarine)</p>	<p>Aquatic ecosystem—high ecological value (HEV)</p>	<p>Ambient (baseflow) Achieve effectively unmodified water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas. Note: there is insufficient information available to establish unmodified water quality for these waters. Refer to QWQG for details on how to establish a minimum water quality data set for deriving local 20th, 50th and 80th percentiles.</p>
<p>Constant Creek subcatchment fresh waters</p>	<p>Aquatic ecosystem—moderately disturbed (MD)</p>	<p>Ambient (baseflow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <10 µg/L^{a, b, c, d} • particulate N: <142 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): <6 µg/L^{a, c} • particulate phosphorus: <20 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 40%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 85%–120% saturation^{a, e, f} • suspended solids: <4 mg/L^a • pH: 7.6–7.9^{a, g, h} • electrical conductivity (EC): <697 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: refer to 'Fresh waters - macroinvertebrates' row (listed after 'Whitsunday islands' in this table) • fish: nd^k

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
		<p>WQOs for pesticides identified in the Mackay Whitsunday WQIP^a:</p> <ul style="list-style-type: none"> • Ametryn: <0.02 µg/L • Atrazine: <LOD • Diuron: <0.07 µg/L • Hexazinone: <0.13 µg/L • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row
<p>Constant Creek subcatchment (end of system)</p>	<p>Aquatic ecosystem—moderately disturbed (MD)</p>	<p>Event (high flow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <300 µg/L^{a, b, d} • particulate N: <279 µg/L^{a, d} • filterable reactive phosphorus (FRP): <30 µg/L^a • particulate phosphorus: <66 µg/L^a • suspended solids: <64 mg/L^a <p>Event WQOs for pesticides^a:</p> <ul style="list-style-type: none"> • Ametryn: <0.05 µg/L • Atrazine: <0.24 µg/L • Diuron: <0.75 µg/L • Hexazinone: <0.29 µg/L • Tebuthiuron: <LOD
<p>Eden Lassie Creek subcatchment fresh waters in area HEV2301</p>	<p>Aquatic ecosystem—high ecological value (HEV)</p>	<p>Ambient (baseflow)</p> <p>Maintain existing water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas.</p> <p>Note: there is insufficient information available to establish current water quality for these waters. Refer to QWQG for details on how to establish a minimum water quality data set for deriving local 20th, 50th and 80th percentiles.</p> <p>WQOs for pesticides identified in the Mackay Whitsunday WQIP, to protect freshwater species at the HEV level of protection^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
<p>Eden Lassie Creek subcatchment fresh waters</p>	<p>Aquatic ecosystem—moderately disturbed (MD)</p>	<p>Ambient (baseflow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <18 µg/L^{a, b, c, d} • particulate N: <39 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): <22 µg/L^{a, c} • particulate phosphorus: <9 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 40%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 85%–120% saturation^{a, e, f} • suspended solids: <1 mg/L^a • pH: 6.5–7.5^{a, g, h} • electrical conductivity (EC): <483 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: nd^k • fish: nd^k <p>WQOs for pesticides identified in the Mackay Whitsunday WQIP^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row
<p>Eden Lassie Creek subcatchment (end of system)</p>	<p>Aquatic ecosystem—moderately disturbed (MD)</p>	<p>Event (high flow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <213 µg/L^{a, b, d} • particulate N: <327 µg/L^{a, d} • filterable reactive phosphorus (FRP): <30 µg/L^a • particulate phosphorus: <70 µg/L^a • suspended solids: <141 mg/L^a <p>Event WQOs for pesticides^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <0.06 µg/L • Hexazinone: <LOD • Tebuthiuron: <LOD
<p>Gregory River subcatchment fresh waters in area HEV2302</p>	<p>Aquatic ecosystem—high ecological value (HEV)</p>	<p>Ambient (baseflow)</p> <p>Maintain existing water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas.</p> <p>The 20th, 50th and 80th percentiles of existing water quality for Gregory River subcatchment waters are:</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: 10–20–31 µg/L^{a, b, c, d} • particulate N: 10–16–52 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): 9–10–15 µg/L^{a, c} • particulate phosphorus: 4–10–17 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 50%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 90%–105% saturation^{a, e, f} • suspended solids: 1–2–3 mg/L^a • pH: 7.2–7.3–7.6^{a, g, h} • electrical conductivity (EC): 180–260–780 µS/cm^{a, i}

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
		<ul style="list-style-type: none"> • temperature: nd^{j, k} • macroinvertebrates: nd^k • fish: nd^k <p>WQOs for pesticides identified in the Mackay Whitsunday WQIP, to protect freshwater species at the HEV level of protection^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row
<p>Gregory River subcatchment fresh waters</p>	<p>Aquatic ecosystem—moderately disturbed (MD)</p>	<p>Ambient (baseflow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <30 µg/L^{a, b, c, d} • particulate N: <43 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): <6 µg/L^{a, c} • particulate phosphorus: <6 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 40%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 85%–120% saturation^{a, e, f} • suspended solids: <2 mg/L^a • pH: 7.2–8.1^{a, g, h} • electrical conductivity (EC): <580 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: refer to 'Fresh waters - macroinvertebrates' row (listed after 'Whitsunday islands' in this table) • fish: nd^k <p>WQOs for pesticides identified in the Mackay Whitsunday WQIP^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row
<p>Gregory River subcatchment (end of system)</p>	<p>Aquatic ecosystem—moderately disturbed (MD)</p>	<p>Event (high flow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <300 µg/L^{a, b, d} • particulate N: <254 µg/L^{a, d} • filterable reactive phosphorus (FRP): <30 µg/L^a • particulate phosphorus: <57 µg/L^a • suspended solids: <42 mg/L^a <p>Event WQOs for pesticides^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <0.06 µg/L • Diuron: <0.31 µg/L • Hexazinone: <0.04 µg/L • Tebuthiuron: <LOD

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
Island freshwaters - Southern Cumberland (refer plans WQ1221, WQ1222)		
Comprising fresh waters on Blacksmith, Brampton, Keswick, Ladysmith, Goldsmith, Hammer, and other southern islands not included in Whitsunday islands		
Southern Cumberland islands - fresh waters in area HEV2385 (various islands)	Aquatic ecosystem—high ecological value (HEV)	<p>Ambient (baseflow)</p> <p>Maintain existing water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas. In the absence of more locally applicable information, the 20th, 50th and 80th percentiles of existing water quality for these island fresh waters are based on the WQOs for Gillinbin Creek subcatchment (Pioneer Basin - mainland):</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: 4–9–13 µg/L^{a, b, c, d} • particulate N: 39–58–152 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): 1–2–3 µg/L^{a, c} • particulate phosphorus: 6–12–22 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 50%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 90%–105% saturation^{a, e, f} • suspended solids: 1–2–4 mg/L^a • pH: 6.7–6.9–7.1^{a, g, h} • electrical conductivity (EC): 130–190–350 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: nd^k • fish: nd^k <p>In the absence of more locally applicable information, WQOs for pesticides are based on those identified in the Mackay Whitsunday WQIP for the Gillinbin Creek subcatchment, to protect freshwater species at the HEV level of protection^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row <p>Refer to QWQG for details on how to establish a minimum water quality data set for deriving local 20th, 50th and 80th percentiles.</p>
Southern Cumberland islands - fresh waters	Aquatic ecosystem—moderately disturbed (MD)	<p>Ambient (baseflow)</p> <p>In the absence of more locally applicable information, the WQOs for these island fresh waters at the moderately disturbed level of protection are based on the 20th/80th percentile of WQOs for Gillinbin Creek subcatchment (Pioneer Basin - mainland):</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <13 µg/L^{a, b, c, d} • particulate N: <152 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): <3 µg/L^{a, c} • particulate phosphorus: <22 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 50%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 90%–105% saturation^{a, e, f} • suspended solids: <4 mg/L^a • pH: 6.7–7.1^{a, g, h} • electrical conductivity (EC): <350 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: nd^k

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
		<ul style="list-style-type: none"> • fish: nd^k <p>In the absence of more locally applicable information, WQOs for pesticides are based on those in the Gillinbin Creek subcatchment^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row <p>Refer to QWQG for details on how to establish a minimum water quality data set for deriving local WQOs.</p>
<p>Island fresh waters - Whitsundays (refer plans WQ1221, WQ1222)</p> <p>Comprising fresh waters on islands north from Waterhole Creek sub catchment boundary near Midge Point through to and including Repulse and Thomas islands, and including islands of the Whitsunday Group (including Dent, Hamilton, Hayman, Hook, Whitsunday islands), the Lindeman Group (including Lindeman, Shaw, and Thomas islands), the Molle group (including Daydream, North and South Molle islands), and the northern group (including Gloucester Island)</p>		
<p>Whitsunday islands -fresh waters in area HEV2384 (various islands)</p>	<p>Aquatic ecosystem—high ecological value (HEV)</p>	<p>Ambient (baseflow)</p> <p>Maintain existing water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas.</p> <p>In the absence of more locally applicable information, the 20th, 50th and 80th percentiles of existing water quality for Whitsunday islands fresh waters are based on the WQOs for Repulse Creek subcatchment (mainland):</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: 10–20–31 µg/L^{a, b, c, d} • particulate N: 10–16–52 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): 9–10–15 µg/L^{a, c} • particulate phosphorus: 4–10–17 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 50%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 90%–105% saturation^{a, e, f} • suspended solids: 1–2–3 mg/L^a • pH: 7.2–7.3–7.6^{a, g, h} • electrical conductivity (EC): 180–260–780 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: nd^k • fish: nd^k <p>In the absence of more locally applicable information, WQOs for pesticides are based on those identified in the Mackay Whitsunday WQIP for the Repulse Creek subcatchment, to protect freshwater species at the HEV level of protection^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row <p>Refer to QWQG for details on how to establish a minimum water quality data set for deriving local 20th, 50th and 80th percentiles.</p>

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
<p>Whitsunday islands fresh waters</p>	<p>Aquatic ecosystem—moderately disturbed (MD)</p>	<p>Ambient (baseflow)</p> <p>In the absence of more locally applicable information, WQOs for Whitsunday islands fresh waters at the moderately disturbed level of protection are based on the 20th/80th percentile of WQOs for Repulse Creek subcatchment (mainland):</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <31 µg/L^{a, b, c, d} • particulate N: <52 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): <15 µg/L^{a, c} • particulate phosphorus: <17 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 50%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 90%–105% saturation^{a, e, f} • suspended solids: <3 mg/L^a • pH: 7.2–7.6^{a, g, h} • electrical conductivity (EC): <780 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: nd^k • fish: nd^k <p>In the absence of more locally applicable information, WQOs for pesticides are based on those identified in the Mackay Whitsunday WQIP for the Repulse Creek subcatchment^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row <p>Refer to QWQG for details on how to establish a minimum water quality data set for deriving local WQOs.</p>
<p>Lethe Brook subcatchment fresh waters in area HEV2306</p>	<p>Aquatic ecosystem—high ecological value (HEV)</p>	<p>Ambient (baseflow)</p> <p>Maintain existing water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas.</p> <p>The 20th, 50th and 80th percentiles of existing water quality for Lethe Brook subcatchment waters are:</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: 9–18–46 µg/L^{a, b, c, d} • particulate N: 21–39–62 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): 12–22–28 µg/L^{a, c} • particulate phosphorus: 4–9–13 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 50%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 90%–105% saturation^{a, e, f} • suspended solids: 0–1–1 mg/L^a • pH: 8.0–8.2–8.3^{a, g, h} • electrical conductivity (EC): 320–430–600 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: nd^k • fish: nd^k <p>WQOs for pesticides identified in the Mackay Whitsunday WQIP, to protect freshwater species at the HEV level of protection^a:</p>

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
		<ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row
Lethe Brook subcatchment fresh waters	Aquatic ecosystem—moderately disturbed (MD)	<p>Ambient (baseflow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <8 µg/L^{a, b, c, d} • particulate N: <101 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): <8 µg/L^{a, c} • particulate phosphorus: <18 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 40%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 85%–120% saturation^{a, e, f} • suspended solids: <3 mg/L^a • pH: 7.5–7.8^{a, g, h} • electrical conductivity (EC): <463 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: refer to 'Fresh waters - macroinvertebrates' row (listed after 'Whitsunday islands' in this table) • fish: nd^k <p>WQOs for pesticides identified in the Mackay Whitsunday WQIP^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <0.01 µg/L • Hexazinone: <0.04 µg/L • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row
Lethe Brook subcatchment (end of system)	Aquatic ecosystem—moderately disturbed (MD)	<p>Event (high flow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <300 µg/L^{a, b, d} • particulate N: <120 µg/L^{a, d} • filterable reactive phosphorus (FRP): <30 µg/L^a • particulate phosphorus: <28 µg/L^a • suspended solids: <38 mg/L^a <p>Event WQOs for pesticides^a:</p> <ul style="list-style-type: none"> • Ametryn: <0.04 µg/L • Atrazine: <0.21 µg/L • Diuron: <0.66 µg/L • Hexazinone: <0.25 µg/L • Tebuthiuron: <LOD

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
<p>Murray Creek subcatchment fresh waters in areas HEV2313 and SD2381 (nb: the majority of SD2381 is estuarine)</p>	<p>Aquatic ecosystem—high ecological value (HEV)</p>	<p>Ambient (baseflow) HEV: Maintain existing water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas. SD: Achieve effectively unmodified water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas. The 20th, 50th and 80th percentiles are:</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: 9–18–46 µg/L^{a, b, c, d} • particulate N: 21–39–62 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): 12–22–28 µg/L^{a, c} • particulate phosphorus: 4–9–13 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 50%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 90%–105% saturation^{a, e, f} • suspended solids: 0–1–1 mg/L^a • pH: 8.0–8.2–8.3^{a, g, h} • electrical conductivity (EC): 320–430–600 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: nd^k • fish: nd^k <p>WQOs for pesticides identified in the Mackay Whitsunday WQIP, to protect freshwater species at the HEV level of protection^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
Murray Creek subcatchment fresh waters	Aquatic ecosystem— moderately disturbed (MD)	<p>Ambient (baseflow)</p> <p>Maintain existing water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas.</p> <p>The 20th, 50th and 80th percentiles of existing water quality for Murray Creek subcatchment waters are:</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: 9–18–46 µg/L^{a, b, c, d} • particulate N: 21–39–62 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): 12–22–28 µg/L^{a, c} • particulate phosphorus: 4–9–13 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 50%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 90%–105% saturation^{a, e, f} • suspended solids: 0–0–1 mg/L^a • pH: 8.0–8.3^{a, g, h} • electrical conductivity (EC): 320–430–600 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: refer to 'Fresh waters - macroinvertebrates' row (listed after 'Whitsunday islands' in this table) • fish: nd^k <p>WQOs for pesticides identified in the Mackay Whitsunday WQIP^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row
Murray Creek subcatchment (end of system)	Aquatic ecosystem— moderately disturbed (MD)	<p>Event (high flow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <300 µg/L^{a, b, d} • particulate N: <206 µg/L^{a, d} • filterable reactive phosphorus (FRP): <30 µg/L^a • particulate phosphorus: <48 µg/L^a • suspended solids: <67 mg/L^a <p>Event WQOs for pesticides^a:</p> <ul style="list-style-type: none"> • Ametryn: <0.05 µg/L • Atrazine: <0.25 µg/L • Diuron: <0.75 µg/L • Hexazinone: <0.3 µg/L • Tebuthiuron: <LOD

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
<p>Myrtle Creek subcatchment fresh waters in areas HEV2307, SD2301</p>	<p>Aquatic ecosystem—high ecological value (HEV)</p>	<p>Ambient (baseflow) HEV: Maintain existing water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas. SD: Achieve effectively unmodified water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas The 20th, 50th and 80th percentiles for Myrtle Creek subcatchment waters are:</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: 10–20–31 µg/L^{a, b, c, d} • particulate N: 10–16–52 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): 9–10–15 µg/L^{a, c} • particulate phosphorus: 4–10–17 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 50%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 90%–105% saturation^{a, e, f} • suspended solids: 1–2–3 mg/L^a • pH: 7.2–7.3–7.6^{a, g, h} • electrical conductivity (EC): 180–260–780 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: nd^k • fish: nd^k • WQOs for pesticides identified in the Mackay Whitsunday WQIP, to protect freshwater species at the HEV level of protection^a: • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row
<p>Myrtle Creek subcatchment fresh waters</p>	<p>Aquatic ecosystem—moderately disturbed (MD)</p>	<p>Ambient (baseflow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <30 µg/L^{a, b, c, d} • particulate N: <112 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): <25 µg/L^{a, c} • particulate phosphorus: <20 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 40%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 85%–120% saturation^{a, e, f} • suspended solids: <5 mg/L^a • pH: 7.2–7.3^{a, g, h} • electrical conductivity (EC): <654 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: refer to 'Fresh waters - macroinvertebrates' row (listed after 'Whitsunday islands' in this table) • fish: nd^k • WQOs for pesticides identified in the Mackay Whitsunday WQIP^a: • Ametryn: <0.04 µg/L • Atrazine: <0.11 µg/L • Diuron: <0.11 µg/L • Hexazinone: <0.08 µg/L • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
Myrtle Creek subcatchment (end of system)	Aquatic ecosystem—moderately disturbed (MD)	<p>Event (high flow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <300 µg/L^{a, b, d} • particulate N: <340 µg/L^{a, d} • filterable reactive phosphorus (FRP): <30 µg/L^a • particulate phosphorus: <70 µg/L^a • suspended solids: <40 mg/L^a <p>Event WQOs for pesticides^a:</p> <ul style="list-style-type: none"> • Ametryn: <0.12 µg/L • Atrazine: <0.94 µg/L • Diuron: <1.5 µg/L • Hexazinone: <0.49 µg/L • Tebuthiuron: <LOD
O'Connell River subcatchment fresh waters in area HEV2310	Aquatic ecosystem—high ecological value (HEV)	<p>Ambient (baseflow)</p> <p>Maintain existing water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas.</p> <p>The 20th, 50th and 80th percentiles of existing water quality for O'Connell River subcatchment waters are:</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: 9–18–46 µg/L^{a, b, c, d} • particulate N: 21–39–62 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): 12–22–28 µg/L^{a, c} • particulate phosphorus: 4–9–13 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 50%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 90%–105% saturation^{a, e, f} • suspended solids: 0–1–1mg/L^a • pH: 8.0–8.2–8.3^{a, g, h} • electrical conductivity (EC): 320–430–600 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: nd^k • fish: nd^k • WQOs for pesticides identified in the Mackay Whitsunday WQIP, to protect freshwater species at the HEV level of protection^a: <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
O'Connell River subcatchment fresh waters	Aquatic ecosystem—moderately disturbed (MD)	<p>Ambient (baseflow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <30 µg/L^{a, b, c, d} • particulate N: <43 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): <6 µg/L^{a, c} • particulate phosphorus: <6 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 40%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 85%–120% saturation^{a, e, f} • suspended solids: <2 mg/L^a • pH: 7.2–8.1^{a, g, h} • electrical conductivity (EC): <580 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: refer to 'Fresh waters - macroinvertebrates' row (listed after 'Whitsunday islands' in this table) • fish: nd^k <p>WQOs for pesticides identified in the Mackay Whitsunday WQIP^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row
O'Connell River subcatchment (end of system)	Aquatic ecosystem—moderately disturbed (MD)	<p>Event (high flow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <300 µg/L^{a, b, d} • particulate N: <340 µg/L^{a, d} • filterable reactive phosphorus (FRP): <30 µg/L^a • particulate phosphorus: <70 µg/L^a • suspended solids: <158 mg/L^a <p>Event WQOs for pesticides^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <0.06 µg/L • Diuron: <0.28 µg/L • Hexazinone: <0.04 µg/L • Tebuthiuron: <0.16 µg/L

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
<p>Proserpine River subcatchment fresh waters in area HEV2305</p>	<p>Aquatic ecosystem—high ecological value (HEV)</p>	<p>Ambient (baseflow) Maintain existing water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas. Note: there is insufficient information available to establish current water quality for these waters. Refer to QWQG for details on how to establish a minimum water quality data set for deriving local 20th, 50th and 80th percentiles. Ambient WQOs for pesticides identified in the Mackay Whitsunday WQIP, to protect freshwater species at the HEV level of protection^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row
<p>Proserpine River subcatchment fresh waters</p>	<p>Aquatic ecosystem—moderately disturbed (MD)</p>	<p>Ambient (baseflow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <30 µg/L^{a, b, c, d} • particulate N: <150 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): <25 µg/L^{a, c} • particulate phosphorus: <20 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 40%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 85%–120% saturation^{a, e, f} • suspended solids: <5 mg/L^a • pH: 6.9–7.5^{a, g, h} • electrical conductivity (EC): <270 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: refer to 'Fresh waters - macroinvertebrates' row (listed after 'Whitsunday islands' in this table) • fish: nd^k <p>WQOs for pesticides identified in the Mackay Whitsunday WQIP^a:</p> <ul style="list-style-type: none"> • Ametryn: <0.04 µg/L • Atrazine: <0.11 µg/L • Diuron: <0.11 µg/L • Hexazinone: <0.08 µg/L • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
Proserpine River subcatchment (end of system)	Aquatic ecosystem—moderately disturbed (MD)	<p>Event (high flow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <300 µg/L^{a, b, d} • particulate N: <340 µg/L^{a, d} • filterable reactive phosphorus (FRP): <30 µg/L^a • particulate phosphorus: <70 µg/L^a • suspended solids: <200 mg/L^a <p>Event WQOs for pesticides^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <0.26 µg/L • Diuron: <1.0 µg/L • Hexazinone: <0.19 µg/L • Tebuthiuron: <0.42 µg/L
Reliance Creek subcatchment (The Leap) fresh waters in area HEV2314	Aquatic ecosystem—high ecological value (HEV)	<p>Ambient (baseflow)</p> <p>Maintain existing water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas.</p> <p>The 20th, 50th and 80th percentiles of existing water quality for Reliance Creek subcatchment waters are:</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: 8–11–17 µg/L^{a, b, c, d} • particulate N: 21–32–81 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): 4–5–9 µg/L^{a, c} • particulate phosphorus: 3–4–5 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – No-flow 20th and 80th percentile: 50%–120% saturation^{a, e, f} – Flow 20th and 80th percentile: 90%–105% saturation^{a, e, f} • suspended solids: 0–1–1 mg/L^a • pH: 7.4–7.6–7.8^{a, g, h} • electrical conductivity (EC): 60–60–70 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: nd^k • fish: nd^k • WQOs for pesticides identified in the Mackay Whitsunday WQIP, to protect freshwater species at the HEV level of protection^a: <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row
Reliance Creek subcatchment (other) freshwaters in areas HEV2315 and SD2381 (nb: the majority of SD2381 is estuarine)	Aquatic ecosystem—high ecological value (HEV)	<p>Ambient (baseflow)</p> <p>HEV: Maintain existing water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas.</p> <p>SD: Achieve effectively unmodified water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas</p> <p>Note: there is insufficient information available to establish water quality objectives for these waters. Refer to QWQG for details on how to establish a minimum water quality data set for deriving local 20th, 50th and 80th percentiles.</p> <p>WQOs for pesticides identified in the Mackay Whitsunday WQIP, to protect freshwater species at the HEV level of protection^a:</p>

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
		<ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row
Reliance Creek subcatchment fresh waters	Aquatic ecosystem—moderately disturbed (MD)	<p>Ambient (baseflow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <30 µg/L^{a, b, c, d} • particulate N: <110 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): <15 µg/L^{a, c} • particulate phosphorus: <20 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 40%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 85%–120% saturation^{a, e, f} • suspended solids: <5 mg/L^a • pH: 7.3–7.5^{a, g, h} • electrical conductivity (EC): <527 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: refer to 'Fresh waters - macroinvertebrates' row (listed after 'Whitsunday islands' in this table) • fish: nd^k <p>WQOs for pesticides identified in the Mackay Whitsunday WQIP^a:</p> <ul style="list-style-type: none"> • Ametryn: <0.02 µg/L • Atrazine: <0.09 µg/L • Diuron: <0.19 µg/L • Hexazinone: <0.20 µg/L • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row
Reliance Creek subcatchment (end of system)	Aquatic ecosystem—moderately disturbed (MD)	<p>Event (high flow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <300 µg/L^{a, b, d} • particulate N: <274 µg/L^{a, d} • filterable reactive phosphorus (FRP): <30 µg/L^a • particulate phosphorus: <70 µg/L^a • suspended solids: <42 mg/L^a <p>Event WQOs for pesticides^a:</p> <ul style="list-style-type: none"> • Ametryn: <0.06 µg/L • Atrazine: <0.61 µg/L • Diuron: <1 µg/L • Hexazinone: <0.41 µg/L • Tebuthiuron: <LOD

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
<p>Repulse Creek subcatchment fresh waters in areas HEV2308 and SD2302</p>	<p>Aquatic ecosystem—high ecological value (HEV)</p>	<p>Ambient (baseflow) HEV: Maintain existing water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas. SD: Achieve effectively unmodified water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas. The 20th, 50th and 80th percentiles for Repulse Creek subcatchment waters are:</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: 10–20–31 µg/L^{a, b, c, d} • particulate N: 10–16–52 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): 9–10–15 µg/L^{a, c} • particulate phosphorus: 4–10–17 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 50%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 90%–105% saturation^{a, e, f} • suspended solids: 1–2–3 mg/L^a • pH: 7.2–7.3–7.6^{a, g, h} • electrical conductivity (EC): 180–260–780 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: nd^k • fish: nd^k <p>WQOs for pesticides identified in the Mackay Whitsunday WQIP, to protect freshwater species at the HEV level of protection^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row
<p>Repulse Creek subcatchment fresh waters</p>	<p>Aquatic ecosystem—moderately disturbed (MD)</p>	<p>Ambient (baseflow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <31 µg/L^{a, b, c, d} • particulate N: <52 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): <15 µg/L^{a, c} • particulate phosphorus: <17 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 50%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 90%–105% saturation^{a, e, f} • suspended solids: <3 mg/L^a • pH: 7.2–7.6^{a, g, h} • electrical conductivity (EC): <780 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: nd^k • fish: nd^k <p>WQOs for pesticides identified in the Mackay Whitsunday WQIP^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
Repulse Creek subcatchment (end of system)	Aquatic ecosystem—high ecological value (HEV)	<p>Event (high flow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <256 µg/L^{a, b, d} • particulate N: <261 µg/L^{a, d} • filterable reactive phosphorus (FRP): <27 µg/L^a • particulate phosphorus: <31 µg/L^a • suspended solids: <8 mg/L^a <p>Event WQOs for pesticides^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD
Southern Cumberland islands fresh waters - refer listing under 'Island fresh waters - Southern Cumberland'		
St Helens Creek subcatchment fresh waters in area HEV2312	Aquatic ecosystem—high ecological value (HEV)	<p>Ambient (baseflow)</p> <p>Maintain existing water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas.</p> <p>The 20th, 50th and 80th percentiles of existing water quality for St Helens Creek subcatchment waters are:</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: 8–11–17 µg/L^{a, b, c, d} • particulate N: 21–32–81 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): 4–5–9 µg/L^{a, c} • particulate phosphorus: 3–4–5 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 50%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 90%–105% saturation^{a, e, f} • suspended solids: 0–1–1 mg/L^a • pH: 7.4–7.6–7.8^{a, g, h} • electrical conductivity (EC): 60–60–70 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: nd^k • fish: nd^k • WQOs for pesticides identified in the Mackay Whitsunday WQIP, to protect freshwater species at the HEV level of protection^a: <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row
St Helens Creek subcatchment fresh waters in area SD2381	Aquatic ecosystem—high ecological value (HEV)	<p>Ambient (baseflow)</p> <p>Achieve effectively unmodified water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas.</p> <p>Note: there is insufficient information available to establish unmodified water quality for these waters. Refer to QWQG for details on how to establish a minimum water quality data set for deriving local 20th, 50th and 80th percentiles.</p>

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
St Helens Creek subcatchment fresh waters	Aquatic ecosystem—moderately disturbed (MD)	<p>Ambient (baseflow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <10 µg/L^{a, b, c, d} • particulate N: <142 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): <6 µg/L^{a, c} • particulate phosphorus: <20 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 40%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 85%–120% saturation^{a, e, f} • suspended solids: <4 mg/L^a • pH: 7.6–7.9^{a, g, h} • electrical conductivity (EC): <697 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: refer to 'Fresh waters - macroinvertebrates' row (listed after 'Whitsunday islands' in this table) • fish: nd^k <p>WQOs for pesticides identified in the Mackay Whitsunday WQIP^a:</p> <ul style="list-style-type: none"> • Ametryn: <0.02 µg/L • Atrazine: <LOD • Diuron: <0.07 µg/L • Hexazinone: <0.13 µg/L • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row
St Helens Creek subcatchment (end of system)	Aquatic ecosystem—moderately disturbed (MD)	<p>Event (high flow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <300 µg/L^{a, b, d} • particulate N: <121 µg/L^{a, d} • filterable reactive phosphorus (FRP): <30 µg/L^a • particulate phosphorus: <33 µg/L^a • suspended solids: <45 mg/L^a <p>Event WQOs for pesticides^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <0.04 µg/L • Diuron: <0.46 µg/L • Hexazinone: <0.23 µg/L • Tebuthiuron: <LOD
Thompson Creek subcatchment fresh waters in area SD2381 (small areas)	Aquatic ecosystem—high ecological value (HEV)	<p>Ambient (baseflow)</p> <p>Achieve effectively unmodified water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas.</p> <p>Note: there is insufficient information available to establish unmodified water quality for these waters. Refer to QWQG for details on how to establish a minimum water quality data set for deriving local 20th, 50th and 80th percentiles.</p>

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
<p>Thompson Creek subcatchment fresh waters</p>	<p>Aquatic ecosystem—moderately disturbed (MD)</p>	<p>Ambient (baseflow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <10 µg/L^{a, b, c, d} • particulate N: <142 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): <6 µg/L^{a, c} • particulate phosphorus: <20 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – No-flow 20th and 80th percentile: 40%–120% saturation^{a, e, f} – Flow 20th and 80th percentile: 85%–120% saturation^{a, e, f} • suspended solids: <4 mg/L^a • pH: 7.6–7.9^{a, g, h} • electrical conductivity (EC): <697 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: refer to 'Fresh waters - macroinvertebrates' row (listed after 'Whitsunday islands' in this table) • fish: nd^k <p>WQOs for pesticides identified in the Mackay Whitsunday WQIP^a:</p> <ul style="list-style-type: none"> • Ametryn: <0.02 µg/L • Atrazine: <LOD • Diuron: <0.07 µg/L • Hexazinone: <0.13 µg/L • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row
<p>Thompson Creek subcatchment (end of system)</p>	<p>Aquatic ecosystem—moderately disturbed (MD)</p>	<p>Event (high flow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <300 µg/L^{a, b, d} • particulate N: <67 µg/L^{a, d} • filterable reactive phosphorus (FRP): <30 µg/L^a • particulate phosphorus: <15 µg/L^a • suspended solids: <22 mg/L^a <p>Event WQOs for pesticides^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <0.15 µg/L • Diuron: <0.46 µg/L • Hexazinone: <0.17 µg/L • Tebuthiuron: <LOD

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
<p>Upper Proserpine River subcatchment fresh waters in area HEV2304</p>	<p>Aquatic ecosystem—high ecological value (HEV)</p>	<p>Ambient (baseflow) Maintain existing water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas. The 20th, 50th and 80th percentiles of existing water quality for Upper Proserpine River subcatchment waters are:</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: 9–18–46 µg/L^{a, b, c, d} • particulate N: 21–39–62 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): 12–22–28 µg/L^{a, c} • particulate phosphorus: 4–9–13 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 50%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 90%–105% saturation^{a, e, f} • suspended solids: 0–1–1mg/L^a • pH: 6.5–8.5^{a, g, h} • electrical conductivity (EC): 320–483–600 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: nd^k • fish: nd^k <p>WQOs for pesticides identified in the Mackay Whitsunday WQIP, to protect freshwater species at the HEV level of protection^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row
<p>Upper Proserpine River subcatchment fresh waters</p>	<p>Aquatic ecosystem—moderately disturbed (MD)</p>	<p>Ambient (baseflow) Maintain existing water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas. The 20th, 50th and 80th percentiles of existing water quality for Upper Proserpine River subcatchment waters are:</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: 9–18–46 µg/L^{a, b, c, d} • particulate N: 21–39–62 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): 12–22–28 µg/L^{a, c} • particulate phosphorus: 4–9–13 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 50%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 90%–105% saturation^{a, e, f} • suspended solids: 0–1–1 mg/L^a • pH: 6.5–8.5^{a, g, h} • electrical conductivity (EC): 320–483–600 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: refer to 'Fresh waters - macroinvertebrates' row (listed after 'Whitsunday islands' in this table) • fish: nd^k <p>WQOs for pesticides identified in the Mackay Whitsunday WQIP^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
		<ul style="list-style-type: none"> • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row
Upper Proserpine River subcatchment (end of system)	Aquatic ecosystem—moderately disturbed (MD)	<p>Event (high flow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <300 µg/L^{a, b, d} • particulate N: <20 µg/L^{a, d} • filterable reactive phosphorus (FRP): <27 µg/L^a • particulate phosphorus: <31 µg/L^a • suspended solids: <10 mg/L^a <p>Event WQOs for pesticides^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD
Waterhole Creek subcatchment fresh waters	Aquatic ecosystem—high ecological value (HEV)	<p>Ambient (baseflow)</p> <p>No HEV waters identified to date. Refer to rows below.</p>
Waterhole Creek subcatchment fresh waters in area SD2381 (nb: the majority of this SD area is estuarine)	Aquatic ecosystem—high ecological value (HEV)	<p>Ambient (baseflow)</p> <p>Achieve effectively unmodified water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas.</p> <p>The 20th, 50th and 80th percentiles of existing water quality for Waterhole Creek subcatchment waters are provided in the following row.</p>
Waterhole Creek subcatchment fresh waters	Aquatic ecosystem—moderately disturbed (MD)	<p>Ambient (baseflow)</p> <p>Maintain existing water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas.</p> <p>The 20th, 50th and 80th percentiles of existing water quality for Waterhole Creek subcatchment waters are:</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: 9–18–46 µg/L^{a, b, c, d} • particulate N: 21–39–62 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): 12–22–28 µg/L^{a, c} • particulate phosphorus: 4–9–13 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 50%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 90%–105% saturation^{a, e, f} • suspended solids: 0–1–1 mg/L^a • pH: 6.5–8.5^{a, g, h} • electrical conductivity (EC): 320–483–600 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: refer to 'Fresh waters - macroinvertebrates' row (listed after 'Whitsunday islands' in this table) • fish: nd^k <p>WQOs for pesticides identified in the Mackay Whitsunday WQIP^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
Waterhole Creek subcatchment (end of system)	Aquatic ecosystem—moderately disturbed (MD)	<p>Event (high flow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <289 µg/L^{a, b, d} • particulate N: <173 µg/L^{a, d} • filterable reactive phosphorus (FRP): <30 µg/L^a • particulate phosphorus: <42 µg/L^a • suspended solids: <74 mg/L^a <p>Event WQOs for pesticides^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <0.03 µg/L • Diuron: <0.11 µg/L • Hexazinone: <0.02 µg/L • Tebuthiuron: <0.04 µg/L
Whitsunday coastal creeks subcatchment fresh waters in area HEV2303	Aquatic ecosystem—high ecological value (HEV)	<p>Ambient (baseflow)</p> <p>Maintain existing water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas.</p> <p>The 20th, 50th and 80th percentiles of existing water quality for Whitsunday Coast subcatchment waters are:</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: 10–20–31 µg/L^{a, b, c, d} • particulate N: 10–16–52 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): 9–10–15 µg/L^{a, c} • particulate phosphorus: 4–10–17 µg/L^{a, c} • dissolved oxygen: <ul style="list-style-type: none"> – no-flow 20th and 80th percentile: 50%–120% saturation^{a, e, f} – flow 20th and 80th percentile: 90%–105% saturation^{a, e, f} • suspended solids: 1–2–3 mg/L^a • pH: 7.2–7.3–7.6^{a, g, h} • electrical conductivity (EC): 180–260–780 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: nd^k • fish: nd^k <p>WQOs for pesticides identified in the Mackay Whitsunday WQIP, to protect freshwater species at the HEV level of protection^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row
Whitsunday coastal creeks subcatchment fresh waters	Aquatic ecosystem—moderately disturbed (MD)	<p>Ambient (baseflow)</p> <p>Maintain existing water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas.</p> <p>The 20th, 50th and 80th percentiles of existing water quality for Whitsunday Coast subcatchment waters are:</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: 10–20–31 µg/L^{a, b, c, d} • particulate N: 10–16–52 µg/L^{a, c, d} • filterable reactive phosphorus (FRP): 9–10–15 µg/L^{a, c} • particulate phosphorus: 4–10–17 µg/L^{a, c} • dissolved oxygen:

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
		<ul style="list-style-type: none"> - no-flow 20th and 80th percentile: 50%–120% saturation^{a, e, f} - flow 20th and 80th percentile: 90%–105% saturation^{a, e, f} • suspended solids: 1–2–3 mg/L^a • pH: 7.2–7.3–7.6^{a, g, h} • electrical conductivity (EC): 180–260–780 µS/cm^{a, i} • temperature: nd^{j, k} • macroinvertebrates: refer to 'Fresh waters - macroinvertebrates' row (listed after 'Whitsunday islands' in this table) • fish: nd^k <p>WQOs for pesticides identified in the Mackay Whitsunday WQIP^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD • other toxicants: refer 'Fresh waters - other toxicants' row
Whitsunday coastal creeks subcatchment (end of system)	Aquatic ecosystem—moderately disturbed (MD)	<p>Event (high flow)</p> <ul style="list-style-type: none"> • dissolved inorganic nitrogen: <256 µg/L^{a, b, d} • particulate N: <261 µg/L^{a, d} • filterable reactive phosphorus (FRP): <27 µg/L^a • particulate phosphorus: <31 µg/L^a • suspended solids: <8 mg/L^a <p>Event WQOs for pesticides^a:</p> <ul style="list-style-type: none"> • Ametryn: <LOD • Atrazine: <LOD • Diuron: <LOD • Hexazinone: <LOD • Tebuthiuron: <LOD
Whitsunday islands fresh waters - refer listing under 'Island fresh waters - Whitsundays'		
FRESH WATERS - MACROINVERTEBRATES		
All HEV waters	Aquatic ecosystem—high ecological value (HEV)	<ul style="list-style-type: none"> • Insufficient data
O'Connell River Basin: fresh water riverine - edge habitat	Aquatic ecosystem—moderately disturbed (MD) ^w	<ul style="list-style-type: none"> • Taxa richness (edge) 20th and 80th percentiles: 20.0–29.0^w • PET richness (edge) 20th and 80th percentiles: 3–5^w • SIGNAL (v2.iv) (edge) 20th and 80th percentiles: 3.53–4.07^w • % sensitive taxa (edge) 20th and 80th percentiles: nd–4.76^w • % tolerant taxa (edge) 20th and 80th percentiles: 35.0–46.67^w
O'Connell River Basin: fresh water riverine - riffle habitat	Aquatic ecosystem—moderately disturbed (MD) ^w	<ul style="list-style-type: none"> • Taxa richness (riffle) 20th and 80th percentiles: 20–26^w • PET richness (riffle) 20th and 80th percentiles: 4.6–7.0^w • SIGNAL (v2.iv) (riffle) 20th and 80th percentiles: 4.36–4.96^w • % sensitive taxa (riffle) 20th and 80th percentiles: 7.37–11.63^w • % tolerant taxa (riffle) 20th and 80th percentiles: 22.74–34.78^w

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
Proserpine River Basin: fresh water riverine - edge habitat	Aquatic ecosystem—moderately disturbed (MD) ^w	<ul style="list-style-type: none"> • Taxa richness (edge) 20th and 80th percentiles: 19–28^w • PET richness (edge) 20th and 80th percentiles: 3–6^w • SIGNAL (v2.iv) (edge) 20th and 80th percentiles: 3.66–4.38^w • % sensitive taxa (edge) 20th and 80th percentiles: nd–6.22^w • % tolerant taxa (edge) 20th and 80th percentiles: 26.18–45.83^w
Proserpine River Basin: fresh water riverine - riffle habitat	Aquatic ecosystem—moderately disturbed (MD) ^w	<ul style="list-style-type: none"> • Taxa richness (riffle) 20th and 80th percentiles: 18–26^w • PET richness (riffle) 20th and 80th percentiles: 5–8^w • SIGNAL (v2.iv) (riffle) 20th and 80th percentiles: 4.57–5.19^w • % sensitive taxa (riffle) 20th and 80th percentiles: 7.69–15.63^w • % tolerant taxa (riffle) 20th and 80th percentiles: 20.0–30.0^w
Whitsunday islands fresh waters	All	<ul style="list-style-type: none"> • Insufficient data
FRESH WATERS - OTHER TOXICANTS (where not listed above)		
Other toxicants for ALL HEV and SD fresh waters in this table	Aquatic ecosystem—high ecological value (HEV)	<p>WQOs are as per AWQG, to protect fresh water species at the HEV level of protection:</p> <ul style="list-style-type: none"> • Toxicants in water^a: refer to AWQG section 3.4—‘water quality guidelines for toxicants’ (including Tables 3.4.1, 3.4.2, and Figure 3.4.1), and AWQG volume 2 (section 8) • Toxicants in sediments: refer to AWQG section 3.5—‘sediment quality guidelines’ (including Table 3.5.1, Figure 3.5.1), and AWQG volume 2 (section 8) <p>Comply with Code of Practice for Antifouling and In-water Hull Cleaning and Maintenance, ANZECC.</p>
Other toxicants for ALL other fresh waters in this table	Aquatic ecosystem—moderately disturbed (MD)	<p>WQOs are as per AWQG, to protect fresh water species at the MD level of protection (identified in AWQG as slightly to moderately disturbed):</p> <ul style="list-style-type: none"> • Toxicants in water^a: refer to AWQG section 3.4—‘water quality guidelines for toxicants’ (including Tables 3.4.1, 3.4.2, and Figure 3.4.1), and AWQG volume 2 (section 8) • Toxicants in sediments: refer to AWQG section 3.5—‘sediment quality guidelines’ (including Table 3.5.1, Figure 3.5.1), and AWQG volume 2 (section 8) <p>Comply with Code of Practice for Antifouling and In-water Hull Cleaning and Maintenance, ANZECC.</p>
Freshwater riparian areas	Aquatic ecosystem—moderately disturbed (MD)	Protect or restore riparian areas. Refer section 3.1.2—riparian WQOs.
Wetlands	Aquatic ecosystem—moderately disturbed (MD)	<p>Objectives as per AWQG and section 3.1.2.</p> <p>Note: for high impact earthworks within Great Barrier Reef wetland protection areas, refer to the guideline 'Protecting Wetlands of High Ecological Significance in Great Barrier Reef Catchments', and the Queensland wetland buffer planning guideline, available from the department's website.</p>

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
GROUNDWATERS		
Groundwaters	Aquatic ecosystem—high ecological value (HEV)	Where groundwaters interact with surface waters, groundwater quality should not compromise identified EVs and WQOs for those waters. Note: the AWQG (2000) recommends that the highest level of protection should be provided to underground aquatic ecosystems, given their high conservation value. Where groundwaters are in good condition the intent is to maintain existing water quality (20th, 50th and 80th percentiles). There is insufficient information available to establish WQOs for this water type ^p . Refer to QWQG for details on how to establish a minimum water quality data set for deriving local 20th, 50th and 80th percentiles.
ESTUARINE AND COASTAL WATERS - listed estuarine, enclosed coastal, open coastal (refer plans WQ1221 and WQ1222) The following WQOs apply to Proserpine River, Whitsunday Island and O'Connell River estuarine and coastal waters within the boundaries shown in plans WQ1221 and WQ1222.		
Estuarine and coastal waters	Aquatic ecosystem—all	Release of sewage from vessels to be controlled in accordance with requirements of the <i>Transport Operations (Marine Pollution) Act 1995</i> and Regulations. (Refer to Maritime Services Queensland website for further information.)
Estuarine waters in areas HEV2303, 2307, 2308, 2314, 2315, 2381, 2384, 2385	Aquatic ecosystem—high ecological value (HEV)	Maintain existing water quality (20th, 50th and 80th percentiles), habitat, biota, flow and riparian areas. Note: there is insufficient information available to establish current water quality for these waters. Refer to QWQG for details on how to establish a minimum water quality data set for deriving local 20th, 50th and 80th percentiles.
Estuarine waters in areas SD2303, 2381	Aquatic ecosystem—high ecological value (HEV)	Achieve effectively unmodified water quality (20th, 50th and 80th percentiles of HEV waters), habitat, biota, flow and riparian areas. Note: there is insufficient information available to establish WQOs for these waters. Refer to QWQG for details on how to establish a minimum water quality data set for deriving local WQOs. <ul style="list-style-type: none"> • toxicants: refer to toxicants row later in this table
Mid-estuary waters (also applies to tidal canals, constructed estuaries within this water type)	Aquatic ecosystem—moderately disturbed (MD)	Note: For waters shown on the plan as being mid estuary (orange colour) and occurring within inter-tidal zone adjacent to the enclosed coastal/lower estuary water type, these waters might have water quality characteristics more in common with the adjacent enclosed coastal/lower estuary water type. Under such circumstances, reference should be made to the WQOs for enclosed coastal/lower estuary water type. <ul style="list-style-type: none"> • ammonia N: <15 µg/L^{c, n} • oxidised N: <30 µg/L^{c, n, s} • dissolved inorganic N: <45 µg/L^{b, c} • organic N: <200 µg/L^{c, n} • total N: <250 µg/L^{c, n} • filterable reactive phosphorus (FRP): <5 µg/L^{c, n} • total P: <20 µg/L^{c, n} • chlorophyll a: <3.0 µg/Lⁿ • dissolved oxygen: 80–105% saturation^{e, n} • suspended solids: nd^{k, n} • turbidity: <10 NTUⁿ • Secchi depth: >1.0 mⁿ • pH: 6.5–8.4ⁿ • temperature: nd^{j, k, n} • toxicants: refer to toxicants row below

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
Toxicants for ALL estuarine waters within this table	All	Toxicants in water and sediment as per AWQG (2000): <ul style="list-style-type: none"> • Toxicants in water^a: refer to AWQG section 3.4—'water quality guidelines for toxicants' (including tables 3.4.1, 3.4.2, and Figure 3.4.1) and AWQG volume 2 (section 8) • Toxicants in sediments: refer to AWQG section 3.5—'sediment quality guidelines' (including Table 3.5.1, Figure 3.5.1) and AWQG volume 2 (section 8) Comply with Code of Practice for Antifouling and In-water Hull Cleaning and Maintenance, ANZECC.
Estuarine riparian areas	Aquatic ecosystem—moderately disturbed (MD)	Protect or restore riparian areas. Refer section 3.1.2—riparian WQOs.
SD2381 enclosed coastal/ lower estuary waters	Aquatic ecosystem—high ecological value (HEV)	Achieve effectively unmodified water quality (20th, 50th and 80th percentiles), habitat and biota. The 20th, 50th and 80th percentiles to be achieved are: <ul style="list-style-type: none"> • ammonia N: 7–10–15 µg/L^{c, n, u} • oxidised N: 2–4–10 µg/L^{c, n, s, u} • total N: 110–120–160 µg/L^{c, n, u} • filterable reactive phosphorus (FRP): 2–3–5 µg/L^{c, n, u} • total P: 10–15–20 µg/L^{c, n, u} • chlorophyll a: 0.8–1.3–2 µg/L^{n, u} • dissolved oxygen: 85–90–105 % saturation^{n, u} • turbidity: nd^v • Secchi: nd^v • suspended solids: nd^{k, n, v} • pH: 8.1–8.3–8.4^{n, u} • temperature: nd^{j, k, n} • toxicants: refer to toxicants rows later in this table
Enclosed coastal/lower estuary waters (including enclosed coastal waters outside marinas, boat harbours, approved spoil grounds, and not identified as HEV or SD)	Aquatic ecosystem—moderately disturbed (MD)	<ul style="list-style-type: none"> • ammonia N: <15 µg/L^{c, n} • oxidised N: <10 µg/L^{c, n, s} • dissolved inorganic N: <25 µg/L^{b, c} • total N: <160 µg/L^{c, n} • filterable reactive phosphorus (FRP): <5 µg/L^{c, n} • total P: <20 µg/L^{c, n} • chlorophyll a: <2.0 µg/Lⁿ • dissolved oxygen: 85–105% saturation^{e, n} • turbidity: <10 NTU^{n, v} • Secchi depth: >1.0 m^{n, v} • suspended solids: nd^{k, n, v} • pH: 8.1–8.4ⁿ • temperature: nd^{j, k, n} • toxicants: refer to toxicants rows later in this table

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
<p>Marinas, boat harbours - including Abel Point Marina, Hamilton Island Marina, Laguna Quays, and Shute Harbour</p>	<p>Aquatic ecosystem—moderately disturbed (MD)</p>	<ul style="list-style-type: none"> • ammonia N: <15 µg/L^{c, n} • oxidised N: <10 µg/L^{c, n, s} • dissolved inorganic N: <25 µg/L^{b, c} • total N: <160 µg/L^{c, n} • filterable reactive phosphorus (FRP): <5 µg/L^{c, n} • total P: <20 µg/L^{c, n} • chlorophyll a: <2.0 µg/Lⁿ • dissolved oxygen: 85–105% saturation^{e, n} • turbidity: <10 NTU^{n, v} • Secchi depth: >1.0 m^{n, v} • suspended solids: nd^{k, n, v} • pH: 8.1–8.4ⁿ • temperature: nd^{i, k, n} • toxicants: refer to toxicants rows later in this table
<p>HEV2381 open coastal waters (Whitsundays - south to Thomas Island) seaward of the plume line^t shown in WQ1222</p>	<p>Aquatic ecosystem—high ecological value (HEV)</p>	<p>Maintain existing water quality (20th, 50th and 80th percentiles), habitat and biota.</p> <p>The 20th, 50th and 80th percentiles to be maintained are:</p> <ul style="list-style-type: none"> • ammonia N: 0–1–9 µg/L^x • oxidised N: 0–1–2 µg/L^x • particulate N: 12–13–15 µg/L^r • total dissolved N: 52–70–95 µg/L^x • total N: 75–95–125 µg/L^x • filterable reactive phosphorus (FRP): 0–1–3 µg/L^x • particulate P: 1.8–2.4–2.8 µg/L^r • total dissolved P: 3–6–9 µg/L^x • total P: 8–11–14 µg/L^x • silicate: 33–49–88 µg/L^x • chlorophyll a: 0.25–0.36–0.54 µg/L^r • dissolved oxygen: 95%–105% saturation^m • turbidity: 0.7–1.1–2.1 NTU^{r, v} • Secchi depth (annual mean): ≥ 10 m^{m, o, v} • suspended solids: 0.9– 1.4– 2.3 mg/L^{r, v} • pH: 8.1–8.3–8.4^m • temperature: <1°C increase above long-term average maximum^o • toxicants: refer to toxicants rows later in this table
<p>HEV2381 enclosed coastal waters (Whitsunday Island)</p>	<p>Aquatic ecosystem—high ecological value (HEV)</p>	<p>Maintain existing water quality (20th, 50th and 80th percentiles), habitat and biota.</p> <p>The 20th, 50th and 80th percentiles to be maintained are:</p> <ul style="list-style-type: none"> • ammonia N: 7–10–15 µg/L^{c, n, u} • oxidised N: 2–4–10 µg/L^{c, n, s, u} • total N: 110–120–160 µg/L^{c, n, u} • filterable reactive phosphorus (FRP): 2–3–5 µg/L^{c, n, u} • total P: 10–15–20 µg/L^{c, n, u} • chlorophyll a: 0.8–1.3–2 µg/L^{n, u} • dissolved oxygen: 85–90–105 % saturation^{n, u} • turbidity: nd^v • Secchi: nd^v • suspended solids: nd^{k, n, v}

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
		<ul style="list-style-type: none"> • pH: 8.1–8.3–8.4^{n, u} • temperature: nd^{j, k, n} • toxicants: refer to toxicants rows later in this table
<p>HEV2382 offshore waters (around Circular Quay Reef), shown in WQ1222</p>	<p>Aquatic ecosystem—high ecological value (HEV)</p>	<p>Maintain existing water quality (20th, 50th and 80th percentiles), habitat and biota.</p> <p>The 20th, 50th and 80th percentiles to be maintained are:</p> <ul style="list-style-type: none"> • ammonia N: 0–2–9 µg/L^x • oxidised N: 0–1–3 µg/L^x • particulate N: 11–14–19 µg/L^x • total dissolved N: 46–63–86 µg/L^x • total N: 70–85–120 µg/L^x • filterable reactive phosphorus (FRP): 0–1–3 µg/L^x • particulate P: 1.4–1.8–2.4 µg/L^x • total dissolved P: 3–5–9 µg/L^x • total P: 6–10–15 µg/L^x • silicate: 18–33–89 µg/L^x • chlorophyll a (annual mean): ≤ 0.45 µg/L^{m, o} • dissolved oxygen: 95%–105% saturation^m • turbidity: <1 NTU^m • Secchi depth (annual mean): ≥ 17m^x • suspended solids: 0.4–0.6–1.1 mg/L^{v, x} • pH: 8.1–8.3–8.4^m • temperature: <1 °C increase above long-term average maximum^o • toxicants: refer to toxicants rows later in this table
<p>HEV2383 open coastal waters (southern Cumberland region - south of Thomas Island), seaward of the plume line^t shown in WQ1222</p>	<p>Aquatic ecosystem—high ecological value (HEV)</p>	<p>Maintain existing water quality (20th, 50th and 80th percentiles), habitat, biota, and flow.</p> <p>The 20th, 50th and 80th percentiles to be maintained are:</p> <ul style="list-style-type: none"> • ammonia N: 1–2–6 µg/L^x • oxidised N: 0–0–1 µg/L^x • particulate N: 14–18–24 µg/L^x • total dissolved N: 44–74–95 µg/L^x • total N: 45–90–115 µg/L^x • filterable reactive phosphorus (FRP): 1–2–4 µg/L^x • particulate P: 1.6–2.1–3.0 µg/L^x • total dissolved P: 4–9–18 µg/L^x • total P: 5–12–21 µg/L^x • silicate: 41–55–96 µg/L^x • chlorophyll a (annual mean): ≤ 0.45 µg/L^{m, o} • dissolved oxygen: 95%–105% saturation^m • turbidity: <1 NTU^m • Secchi depth (annual mean): ≥ 10 m^{m, o, v} • suspended solids: 1.1–1.6–2.4 mg/L^{r, v} • pH: 8.1–8.3–8.4^m • temperature: <1 °C increase above long-term average maximum^o • toxicants: refer to toxicants rows later in this table

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
<p>SD2381 open coastal waters, landward of the plume line[†] shown in WQ1222</p>	<p>Aquatic ecosystem—high ecological value (HEV)</p>	<p>Achieve effectively unmodified water quality (20th, 50th and 80th percentiles), habitat, biota, and flow.</p> <p>The 20th, 50th and 80th percentiles to be achieved are:</p> <ul style="list-style-type: none"> • ammonia N: 0–1–9 µg/L^x • oxidised N: 0–1–2 µg/L^x • particulate N: 12–13–15 µg/L^r • total dissolved N: 52–70–95 µg/L^x • total N: 75–95–125 µg/L^x • filterable reactive phosphorus (FRP): 0–1–3 µg/L^x • particulate P: 1.8–2.4–2.8 µg/L^r • total dissolved P: 3–6–9 µg/L^x • total P: 8–11–14 µg/L^x • silicate: 33–49–88 µg/L^x • chlorophyll a: 0.25–0.36–0.54 µg/L^r • dissolved oxygen: 95%–105% saturation^m • turbidity: 0.7–1.1–2.1 NTU^{r, v} • Secchi depth (annual mean): ≥ 10 m^{m, o, v} • suspended solids: 0.9–1.4– 2.3 mg/L^{r, v} • pH: 8.1–8.4^m • temperature: <1 °C increase above long-term average maximum^o • toxicants: refer to toxicants rows later in this table
<p>Open coastal waters landward of the plume line[†], shown in WQ1222</p> <p>(including open coastal waters outside marinas, boat harbours, and not identified as HEV or SD)</p>	<p>Aquatic ecosystem—slightly to moderately disturbed (SMD), mapped as MD</p>	<ul style="list-style-type: none"> • particulate N (annual mean): ≤ 20µg/L^{m, o} • particulate P (annual mean): ≤ 2.8 µg/L^{m, o} • chlorophyll a (annual mean): ≤ 0.45 µg/L^{m, o} • dissolved oxygen: 95–105 % saturation^m • turbidity (median): <1 NTU^{m, v} • Secchi depth (annual mean): ≥ 10 m^{m, o, v} • suspended solids (annual mean): ≤ 2.0 mg/L^{m, o, v} • pH: 8.1–8.4^m • temperature: <1 °C increase above long-term average maximum^o • toxicants: refer to toxicants rows later in this table • Note: there is insufficient information available to establish WQOs for the following indicators in these waters: • ammonia N, oxidised N, dissolved inorganic N, total N, filterable reactive phosphorus, total P. • Refer to QWQG for details on how to establish a minimum water quality data set for deriving local WQOs.
COASTAL WATERS - TOXICANTS		
<p>Toxicants for ALL enclosed and open coastal waters in this table, excluding waters in marinas, boat harbours, approved spoil grounds, tidal canals, constructed estuaries</p>	<p>Aquatic ecosystem—high ecological value (HEV)</p>	<p>WQOs for pesticides identified in the Mackay-Whitsunday WQIP, to protect marine species at the HEV level of protection^o:</p> <ul style="list-style-type: none"> • Ametryn: <0.5 µg/L • Atrazine: <0.6 µg/L • Diuron: <0.9 µg/L • Hexazinone: <1.2 µg/L • Tebuthiuron: <0.02 µg/L • Tributyltin: <0.0004 µg/L <p>WQOs for all other toxicants in these waters as per GBRMPA and AWQG water quality guidelines, to protect marine species at the HEV level of protection. For toxicants not listed in GBRMPA guidelines:</p>

Water area/type (refer plans WQ1221, WQ1222)	Management intent (level of protection)	Water quality objectives to protect aquatic ecosystem EV
		<ul style="list-style-type: none"> • Toxicants in water^a: refer to AWQG section 3.4—'water quality guidelines for toxicants' (including tables 3.4.1, 3.4.2, and Figure 3.4.1), and AWQG volume 2 (section 8) • Toxicants in sediments: refer to AWQG section 3.5—'sediment quality guidelines' (including Table 3.5.1, Figure 3.5.1), and AWQG volume 2 (section 8) • Comply with Code of Practice for Antifouling and In-water Hull Cleaning and Maintenance, ANZECC.
<p>Toxicants for marinas, boat harbours, approved spoil grounds, tidal canals, constructed estuaries</p>	<p>Aquatic ecosystem—moderately disturbed (MD) (pesticides: high ecological value (HEV))</p>	<p>WQOs for pesticides identified in the Mackay-Whitsunday WQIP, to protect marine species at the HEV level of protection^o:</p> <ul style="list-style-type: none"> • Ametryn: <0.5 µg/L • Atrazine: <0.6 µg/L • Diuron: <0.9 µg/L • Hexazinone: <1.2 µg/L • Tebuthiuron: <0.02 µg/L • Tributyltin: <0.006 µg/L (SMD / 95% species protection) <p>WQOs for all other toxicants in these waters as per GBRMPA and AWQG water quality guidelines, to protect marine species at the MD level of protection (identified in the AWQG as slightly to moderately disturbed). For toxicants not listed in GBRMPA guidelines:</p> <ul style="list-style-type: none"> • Toxicants in water^a: refer to AWQG section 3.4—'water quality guidelines for toxicants' (including Tables 3.4.1, 3.4.2, and Figure 3.4.1), and AWQG volume 2 (section 8) • Toxicants in sediments: refer to AWQG section 3.5—'sediment quality guidelines' (including Table 3.5.1, Figure 3.5.1), and AWQG volume 2 (section 8) <p>Comply with Code of Practice for Antifouling and In-water Hull Cleaning and Maintenance, ANZECC.</p>
WATER QUALITY REFERENCE SITES		
<p>Fresh, estuarine, and coastal waters: water quality reference (least disturbed) sites</p>	<p>Aquatic ecosystem—high ecological value (HEV)</p>	<p>For sites identified in the Queensland Water Quality Guidelines, the water quality improvement plan, or by other recognised entities under the EPP Water as reference (least disturbed) sites for water quality monitoring, the management intent is to maintain or achieve effectively unmodified water quality (20th, 50th and 80th percentiles), habitat, biota, and flow.</p>

Notes:

Units:

µg/L = micrograms/Litre; mg/L = milligrams/Litre; µS/cm = microSiemens/centimetre; mg/kg = milligrams/kilogram

LOD = limit of detection, which, in fresh waters, is currently 0.01 µg/L for all herbicides in this table.

- The values for these indicators are based on the QWQG Mackay Whitsunday sub-regional water quality guidelines, which are derived from values in the Mackay Whitsunday Water Quality Improvement Plan (WQIP). These include objectives for both ambient/baseflow conditions and high flow ('event') conditions. The event quality objectives are expressed as end of system (e.g. subcatchment) event mean concentrations (EMC). Further details are provided in source documents (refer below).
- DIN = ammonia N + oxidised N.
- Ambient (baseflow) nutrient objectives do not apply during high flow events. Refer to 'event/high flow' rows for applicable high flow objectives. See QWQG Section 5 and Appendix D for more information on applying guidelines under high flow conditions.
- During periods of low flow and particularly in smaller creeks, build up of organic matter derived from natural sources (e.g. leaf litter) can result in increased organic N levels (generally in the range of 400 to 800µg/L). This may lead to total N values exceeding the WQOs. Provided that levels of inorganic N (i.e. NH₃ + oxidised N) remain low, then the elevated levels of organic N should not be seen as a breach of the WQOs, provided this is due to natural causes.
- Dissolved oxygen (DO) objectives apply to daytime conditions. Lower values will occur at night in most waters. In estuaries, reductions should only be in the region of 10–15 per cent saturation below daytime values. In freshwaters, night time reductions are more variable. Following significant rainfall events, reduced DO values may occur due to the influx of organic material. In estuaries post-event values as low as 40 per cent saturation may occur naturally for short periods but values well below this would indicate some anthropogenic effect. In freshwaters, post-event DO reductions are again more variable. In general, DO values consistently less than 50 per cent are likely to impact on the ongoing ability of fish to persist in a waterbody while short term DO values less than 30 per cent saturation are toxic to some

- fish species. Very high DO (supersaturation) values can be toxic to some fish as they cause gas bubble disease.
- f) DO values for fresh waters should only be applied to flowing waters. Stagnant pools in intermittent streams naturally experience values of DO below 50 per cent saturation.
- g) Wallum/tannin-stained waters contain naturally high levels of humic acids (and have a characteristic brown tea-tree stain). In these types of waters, natural pH values may range from 3.6 to 6.
- h) During flood events or nil flow periods, pH values should not fall below 5.5 (except in wallum/tannin waters) or exceed 9.
- i) Conductivity, under natural conditions, is highly dependent on local geology and soil types. Values in this table are based on QWQG Mackay Whitsunday sub regional water quality guidelines. In the absence of sub-regional conductivity WQOs, the QWQG (Appendix G) provides information on conductivity values in a set of 18 defined salinity zones throughout Queensland. For each zone, the QWQG provide a range of percentile values based on data from all the sites within that zone. This provides a useful first estimate of background conductivity within a zone. However, even within zones there is a degree of variation between streams and therefore the values for the zone would still need to be ground truthed against local values.
- j) Temperature varies both daily and seasonally, it is depth dependent and is also highly site specific. It is therefore not possible to provide simple generic WQOs for this indicator. The recommended approach is that local WQOs be developed. Thus, WQOs for potentially impacted streams should be based on measurements from nearby streams that have similar morphology and which are thought not to be impacted by anthropogenic thermal influences. From an ecological effects perspective, the most important aspects of temperature are the daily maximum temperature and the daily variation in temperature. Therefore measurements of temperature should be designed to collect information on these indicators of temperature and, similarly, local WQOs should be expressed in terms of these indicators. Clearly, there will be an annual cycle in the values of these indicators and therefore a full seasonal cycle of measurements is required to develop guideline values.
- k) nd = insufficient or no data, n/a = not applicable for this indicator and water type, ng = no guideline. Will be updated if guidelines become available.
- l) For Andromache, Murray, Upper Proserpine, Waterhole, and Whitsunday Coast, current water quality was comparable to adjacent HEV waters. The WQOs for these waters have been set to maintain current water quality (based on QWQG sub-regional guidelines and WQIP).
- m) The values for these indicators are based on the QWQG Central Coast regional water quality guidelines.
- n) The values for these indicators are based on the QWQG Wet Tropics regional water quality guidelines.
- o) The values for these indicators are based on GBRMPA (2010) Water Quality Guidelines for the Great Barrier Reef Marine Park 2010 (refer 'sources' below). Refer to GBRMPA Water Quality Guidelines for further details. WQOs for coastal water pesticides in this table are based on GBRMPA species protection levels. For Hexazinone there is insufficient data to determine species protection levels. Note that coastal water values for Diazinon, Hexazinone, and Tebuthiuron are based on low reliability guidelines that may be updated with additional information. Refer to GBRMPA Water Quality Guidelines for further details. Also refer to note 'q' re toxicants.
- p) Local groundwater WQOs have not been developed for this region.
- q) The values for these indicators are based on the AWQG. Under the AWQG, a single set of values is provided for the 'slightly to moderately disturbed' (SMD) level of protection, which is identified in this document as 'moderately disturbed' (MD). The HEV and SMD levels of protection typically correspond to 99 per cent and 95 per cent species protection, respectively. For a small number of toxicants with potential toxicity and bioaccumulation effects, the AWQG identify SMD protection level values corresponding to 99% species protection. For further details on toxicant guidelines, e.g. toxicant species, and variability in relation to water quality characteristics (such as pH, hardness) refer to AWQG.
- r) The values for these indicators are based on monitoring data and analysis by GBRMPA. For particulate N and P and SS, values are based on water quality monitoring data (dry and wet season) at Double Cone Island (2005 to 2011; Schaffelke et al, 2011). Chl-a values are based on grouped data from Hook Passage, Cid, Henning, Dent Passage and west of Lindeman Island (1996 to 2009). Turbidity is based on logger data at Double Cone Island (2007 to 2011).
- s) Oxidised N = NO₂ + NO₃. Units for nitrogen indicators are micrograms per litre (µg/L) N.
- t) The GBR plume discharge area is derived from combining the high and very high frequency influence areas of river discharges that contain high and very high pollutant loads including sediment and nutrient. Refer to Devlin *et al* in 'sources' below for more details.
- u) The values for these indicators are based on monitoring data and analysis by the department.
- v) Suspended solids (and hence turbidity and Secchi depth) levels in coastal waters are naturally highly variable depending on wind speed/wave height and in some cases on tidal cycles. The values in this table provide guidance on what the long term values of turbidity, Secchi depth or TSS should comply with. However, these values will often be naturally exceeded in the short term during windy weather or spring tides. They therefore should not be used for comparison with short term data sets. Where assessable coastal developments are proposed, proponents should carry out site specific intensive monitoring of these indicators (or equivalent light penetration indicators) and use these as a baseline for deriving local guidelines and for comparison with post development conditions.
- w) Macroinvertebrates: WQOs for macroinvertebrates are preliminary. The median value of macroinvertebrate indicators at test sites is to be compared and assessed against the macroinvertebrate values in this table. The values do not apply to waters identified for HEV level of protection/management intent. Values were derived for the 'slightly to moderately disturbed' (SMD) level of protection, which is identified in this document as 'moderately disturbed' (MD). Refer to sources below for monitoring/assessment protocols.
Habitats (i.e. edge, riffle): Edge habitat is located along the stream bank. Riffle habitat is characterised as in a reach with relatively steep, shallow (<0.3 m), fast flowing (>0.2 m/s) and broken water over stony beds.
Sample processing: Samples are identified in the laboratory to family level, except for Chironomidae that are identified to sub-family, and lower Phyla (Porifera, Nematoda, Nemertea, etc.), Oligochaeta (freshwater worms), Acarina (mites), and microcrustacea (Ostracoda, Copepoda, Cladocera) that are not identified further. The taxonomy of macroinvertebrates can change over time. The taxonomy used to calculate these indices are based on those used in determining the SIGNAL 2.iv taxa scores.
Taxa richness: Taxa richness is the number of aquatic macroinvertebrate taxa collected in a sample. Use of taxa richness is based on the premise that with changes in the condition of a site the taxa richness will increase or decrease from 'reference'. Increases or decreases will depend on the nature of the threats that are influencing the ecosystem.
PET taxa richness: PET taxa richness (or EPT) is the number of aquatic macroinvertebrate families collected from three orders of aquatic insects: Ephemeroptera (mayflies), Plecoptera (stoneflies) and Trichoptera (caddisflies). Macroinvertebrates belonging to these three

orders are considered to be sensitive to changes in their environment and therefore PET can be used to assess condition.

SIGNAL Index: The Stream Invertebrate Grade Number - Average Level (SIGNAL) index was developed for the bioassessment of water quality in rivers in Australia. It is calculated by grading each macroinvertebrate family based upon the level of its sensitivity to various pollutants. The grades applied range from 1 (tolerant) to 10 (sensitive) (Chessman, 2003). The SIGNAL score for a sample is calculated by averaging the sensitivity grades of the macroinvertebrate families collected. The version applied here is SIGNAL version 2.iv (Chessman, 2003).

% tolerant taxa, % sensitive taxa: are indices based on the proportion of taxa with a SIGNAL grade that are determined to have 'tolerant' sensitivity grades (SIGNAL grades 1, 2 and 3) and 'sensitive' sensitivity grades (SIGNAL grades 8, 9 and 10) respectively. A change in these may reflect changes in water quality, habitat condition, or other causative factors influencing these indices. The version of SIGNAL grades applied here is SIGNAL version 2.iv.

- x) The values for these indicators are based on monitoring conducted from 1985 to 2006, supporting publication by De'ath and Fabricius (2008). Refer to sources below.

Sources:

The following sources contain further background information relevant to WQOs:

ANZECC (1997) Code of Practice for Antifouling and In-water Hull Cleaning and Maintenance.

ANZECC & ARMCANZ (2000) Australian and New Zealand Guidelines for Fresh and Marine Water Quality (AWQG).

Chessman, B (2003) SIGNAL 2 – A Scoring System for Macro-invertebrate ('Water Bugs') in Australian Rivers, Monitoring River Health Initiative Technical Report number 31, Commonwealth of Australia, Canberra.

De'ath G, Fabricius KE (2008) Water quality of the Great Barrier Reef: distributions, effects on reef biota and trigger values for the conservation of ecosystem health. Research publication no. 89. Great Barrier Marine Park Authority, Report to the Great Barrier Reef Marine Park Authority and published by the Great Barrier Reef Marine Park Authority, Townsville, 104 pp

Devlin, M, Harkness, P, McKinna, L & Waterhouse, J (2011) Mapping the surface exposure of terrestrial pollutants in the Great Barrier Reef. Report to the Great Barrier Reef Marine Park Authority, August 2010. Australian Centre for Tropical Freshwater Research. Report Number 10/12.

Drewry, J, Higham, W & Mitchell, C (2008) Water quality improvement plan (WQIP). Final report for the Mackay Whitsunday region. Mackay Whitsunday Natural Resource Management Group.

Drewry, J, Higham, W, Mitchell, C, Rohde, K, Masters, B & Galea, L (2008) Water quality improvement plan (WQIP). Turning environmental values into water quality objectives and targets. Mackay Whitsunday Natural Resource Management Group.

Drewry, J, Higham, W, Mitchell, C & Rohde, K (2008) Water quality improvement plan (WQIP). Modelling sediment and nutrient exports and management scenarios. Mackay Whitsunday Natural Resource Management Group.

Galea, L, Peplinkhouse, D, Loft, F & Folkers, A (2008) Mackay Whitsunday Healthy Waterways Baseline Monitoring Program Regional Report. Queensland Department of Natural Resources and Water for the Mackay Whitsunday Natural Resource Management Group, Australia.

Great Barrier Reef Marine Park Authority (2010) Water quality guidelines for the Great Barrier Reef Marine Park 2010, Great Barrier Reef Marine Park Authority, Townsville, available on the Great Barrier Reef Marine Park Authority's website.

Negus P, Steward A and Blessing J (2013). Macroinvertebrate water quality guidelines: Townsville and Mackay-Whitsunday regions. Water Planning Ecology Group, Science Delivery Division. DSITIA, The State of Queensland.

NRM (2001). Australia-wide Assessment of River Health: Queensland AusRivAS Sampling and Processing Manual. Monitoring River Health Initiative Technical Report Number 12. Environment Australia, Canberra.

Queensland Government (2009, as amended) Queensland Water Quality Guidelines. (Refer to section 5 and Appendix D of the QWQG for more detail on compliance assessment protocols.)

Rohde, K, Masters, B, Fries, N, Noble, R & Carroll, C (2008) Fresh and Marine Water Quality in the Mackay Whitsunday Region 2004/05 to 2006/07. Queensland Department of Natural Resources and Water for the Mackay Whitsunday Natural Resource Management Group, Australia.

Schaffelke B, Carleton J, Doyle J, Furnas M, Gunn K, Skuza M, Wright M, Zagorskis I (2011) Reef Rescue Marine Monitoring Program. Final Report of AIMS Activities 2010/11– Inshore Water Quality Monitoring. Report for the Great Barrier Reef Marine Park Authority. Australian Institute of Marine Science, Townsville. (83 p.). Additional years also published accessible for download from GBRMPA.

Transport Operations (Marine Pollution) Act 1995 and Regulations 2008, available on the Office of Queensland Parliamentary Counsel website.

3.1.2 Riparian water quality objectives

For vegetation management relating to waterways, reference should be made to the relevant regional vegetation management codes under the *Vegetation Management Act 1999*. These codes include performance requirements relating to watercourses and wetlands, aimed at maintaining water quality, bank stability, aquatic and terrestrial habitat. Codes include vegetation clearing controls that vary according to stream order.

To review the latest applicable vegetation management code (and other explanatory information) for waters for this area, contact the Department of Natural Resources and Mines website.

Planning schemes under the *Sustainable Planning Act 2009* may also specify riparian buffers (for example under catchment protection or waterway codes). Contact the Department of State Development, Infrastructure and Planning website and local government websites for further information about planning schemes.

3.1.3 Stormwater management design objectives

Stormwater management design objectives for urban development are detailed in the department's Urban Stormwater Quality Planning Guidelines 2010 (as amended). Stormwater quality and flow management design objectives are specified for both the construction and operational phases of development in accordance with landscape features and the regional location of proposed development. The guidelines are available on the department's website.

3.2 Water quality objectives for human use environmental values

This section outlines WQOs to protect human use EVs, which comprise those EVs other than the aquatic ecosystem EV (e.g. recreation, stock watering, aquaculture and crop irrigation). Table 1 of this document outlines the EVs that have been identified for different waters in the catchment. Where a human use EV has been identified, the following tables can be used to identify the WQOs to support that EV. Where Table 1 indicates more than one EV applies to a given water (for example aquatic ecosystem and recreational use), the adoption of the most stringent WQO for each water quality indicator will then protect all identified EVs.

WQOs in this section are, unless otherwise specified, based on relevant national water quality guidelines including AWQG and the ADWG⁶. Table 3 outlines human use EVs, applicable water types, and a selection of more commonly used WQOs to support those EVs. Tables 4 to 12 provide further WQOs to protect particular human use EVs (based on national guidelines or other more local studies). Where national guidelines or other codes remain the primary source for WQOs, reference to those national guidelines or codes is necessary to obtain comprehensive listings of all indicators and corresponding WQOs.

Table 3. Water quality objectives to protect human use environmental values.

Environmental value	Water type/area (refer Table 1 and plans WQ1221, WQ1222)	Water quality objectives to protect EV (refer to specified codes and guidelines for full details)
Suitability for drinking water supply	All fresh waters including groundwaters	Local WQOs for drinking water supply are provided in Table 4. Note: For water quality after treatment or at point of use refer to legislation and guidelines, including: <ul style="list-style-type: none"> • <i>Public Health Act 2005</i> and Regulations • <i>Water Supply (Safety and Reliability) Act 2008</i>, including any approved drinking water quality management plan under the Act • <i>Water Fluoridation Act 2008</i> • ADWG 2011.
Protection of the human consumer for oystering	Estuarine and coastal waters	Objectives as per AWQG and Australia New Zealand Food Standards Code ⁷ , Food Standards Australia New Zealand, 2007 and updates.
Protection of the human consumer	Fresh waters, estuarine and coastal waters	Objectives as per AWQG and Australia New Zealand Food Standards Code, Food Standards Australia New Zealand, 2007 and updates.
Protection of cultural and spiritual values	Fresh waters (including groundwaters), estuarine and coastal waters	Protect or restore indigenous and non-indigenous cultural heritage consistent with relevant policies and plans.
Suitability for industrial use	Fresh waters, estuarine and coastal waters	No WQOs are provided in this scheduling document for industrial uses. Water quality requirements for industry vary within and between industries. The AWQG do not provide guidelines to protect industries, and indicate that industrial water quality requirements need to be considered on a case-by-case basis. This EV is usually protected by other values, such as the aquatic ecosystem EV.

⁶ The AWQG are available on the National Water Quality Management Strategy website.

The ADWG are available on the NHMRC website.

⁷ The Australia New Zealand Food Standards Code is available on the Food Standards Australia and New Zealand website.

Environmental value	Water type/area (refer Table 1 and plans WQ1221, WQ1222)	Water quality objectives to protect EV (refer to specified codes and guidelines for full details)
Suitability for aquaculture	Fresh waters, estuarine and coastal waters	Objectives as per: <ul style="list-style-type: none"> • tables 5–7 • AWQG and Australia New Zealand Food Standards Code, Food Standards Australia New Zealand, 2007 and updates.
Suitability for irrigation	All fresh waters including groundwaters	ANZECC objectives for pathogens and metals are provided in tables 8 and 9. For other indicators, such as salinity, sodicity and herbicides, see AWQG.
Suitability for stock watering	All fresh waters including groundwaters	Objectives as per AWQG, including median faecal coliforms <100 organisms per 100 mL. WQOs for total dissolved solids and metals are provided in tables 10 and 11, based on AWQG. For other objectives, such as cyanobacteria and pathogens, see AWQG.
Suitability for farm supply/use	All fresh waters including groundwaters	Objectives as per AWQG.
Suitability for primary contact recreation	Fresh waters, estuarine and coastal waters	Objectives as per NHMRC (2008) ⁸ , including: <ul style="list-style-type: none"> • water free of physical (floating and submerged) hazards • temperature range: 16–34°C • pH range: 6.5–8.5 • DO: >80% • faecal contamination: designated recreational waters are protected against direct contamination with fresh faecal material, particularly of human or domesticated animal origin. Two principal components are required for assessing faecal contamination: <ul style="list-style-type: none"> – assessment of evidence for the likely influence of faecal material – counts of suitable faecal indicator bacteria (usually <i>enterococci</i>) These two components are combined to produce an overall microbial classification of the recreational water body: • intestinal enterococci: 95th percentile ≤ 40 organisms per 100mL (for healthy adults) (NHMRC, 2008; Table 5.7) • direct contact with venomous or dangerous aquatic organisms should be avoided. Recreational water bodies should be reasonably free of, or protected from, venomous organisms (e.g. box jellyfish and bluebottles) • waters contaminated with chemicals that are either toxic or irritating to the skin or mucous membranes are unsuitable for recreational purposes.
Suitability for primary contact recreation	Fresh waters	<ul style="list-style-type: none"> • cyanobacteria/algae: Recreational water bodies should not contain: <ul style="list-style-type: none"> ○ level 1¹: ≥ 10 µg/L total microcystins; or ≥ 50 000 cells/mL toxic <i>Microcystis aeruginosa</i>; or biovolume equivalent of ≥ 4 mm³/L for the combined total of all cyanobacteria where a known toxin producer is dominant in the total biovolume or ○ level 2¹: ≥ 10 mm³/L for total biovolume of all cyanobacterial material where known toxins are not present ○ cyanobacterial scums consistently present. Further details are contained in NHMRC (2008) and Table 12.
	Estuarine, coastal waters	<ul style="list-style-type: none"> • cyanobacteria/algae: Recreational water bodies should not contain ≥ 10 cells/mL <i>Karenia brevis</i> and/or have <i>Lyngbya majuscula</i> and/or <i>Pfiesteria</i> present in high numbers². Further details are contained in NHMRC (2008) and Table 12.

⁸ Guidelines for Managing Risks in Recreational Water are available on the NHMRC website.

Environmental value	Water type/area (refer Table 1 and plans WQ1221, WQ1222)	Water quality objectives to protect EV (refer to specified codes and guidelines for full details)
Suitability for secondary contact recreation	Fresh waters, estuarine and coastal waters	Objectives as per NHMRC (2008), including: <ul style="list-style-type: none"> • intestinal enterococci: 95th percentile ≤ 40 organisms per 100mL (for healthy adults) (NHMRC, 2008; Table 5.7) • cyanobacteria/algae—refer objectives for primary recreation, NHMRC (2008) and Table 12.
Suitability for visual recreation	Fresh waters, estuarine and coastal waters	Objectives as per NHMRC (2008), including: <ul style="list-style-type: none"> • recreational water bodies should be aesthetically acceptable to recreational users. The water should be free from visible materials that may settle to form objectionable deposits; floating debris, oil, scum and other matter; substances producing objectionable colour, odour, taste or turbidity; and substances and conditions that produce undesirable aquatic life. • cyanobacteria/algae—refer objectives for primary recreation, NHMRC (2008) and Table 12.

Notes:

1. Level 1 recognises the probability of adverse health effects from ingestion of known toxins, in this case based on the toxicity of microcystins. Level 2 covers circumstances in which there are very high cell densities of cyanobacterial material, irrespective of the presence of toxicity or known toxins. Increased cyanobacterial densities increase the likelihood of non-specific adverse health outcomes, principally respiratory, irritation and allergy symptoms. (NHMRC, 2008; 8).

2. The NHMRC states that its guidelines are concerned *'only with risks that may be associated with recreational activities in or near coastal and estuarine waters. This includes exposure through dermal contact, inhalation of sea-spray aerosols and possible ingestion of water or algal scums, but does not include dietary exposure to marine algal toxins.'* (NHMRC, 2008; 121).

Sources:

The WQOs were determined from a combination of documents, including:

Australian Drinking Water Guidelines (NHMRC, 2011).

Australia New Zealand Food Standards Code (Australian Government: Food Standards Australia New Zealand).

Australian and New Zealand Guidelines for Fresh and Marine Water Quality (ANZECC & ARMCANZ, 2000).

Guidelines for Managing Risks in Recreational Water (NHMRC, 2008).

Table 4. Drinking water EV: Priority water quality objectives for drinking water supply in the vicinity of off-takes, including groundwater, before treatment.

This table outlines WQOs for water **before treatment**, unless otherwise stated. For water quality after treatment or at the point of use, refer to relevant legislation and guidelines, including *Public Health Act 2005* and Regulations, *Water Supply (Safety and Reliability) Act 2008*, including any approved drinking water quality management plan under the Act, *Water Fluoridation Act 2008*, and the Australian Drinking Water Guidelines (ADWG, 2011). Objectives are derived following advice from Mackay Water, Whitsunday Regional Council, Queensland Health and the Queensland Water Supply Regulator.

Indicator	Water quality objective ¹
<i>Giardia</i>	0 cysts (Queensland Water Supply Regulator) If <i>Giardia</i> is detected in drinking water then the health authorities should be notified immediately and an investigation of the likely source of contamination undertaken (ADWG).
<i>Cryptosporidium</i>	0 cysts (Queensland Water Supply Regulator) If <i>Cryptosporidium</i> is detected in drinking water then the health authorities should be notified immediately and an investigation of the likely source of contamination undertaken (ADWG).
<i>E. coli</i>	<100 cfu/100mL
Enterococci	<100 cfu/100mL Often in large catchments Enterococci is a better indicator of faecal contamination since it is shown to live longer in the environment
Blue-green algae (cyanobacteria)	<10000 cells/mL
Algal toxin	<1 µg/L Microcystin
Turbidity	<30 NTU
Colour	<35 TCU
pH	6.5–8.0
Total hardness	<115 mg/L
Conductivity	<300 µS/cm
Total dissolved solids	ADWG 2011 aesthetic guideline: <600 mg/L
Total organic carbon	<2 mg/L
Sodium	Raw water supply: <25 mg/L General ² : The concentration of sodium in reticulated drinking water supplies should not exceed 180 mg/L (ADWG, based on threshold at which taste becomes appreciable). At-risk groups (medical) ² : The concentration of sodium in water supplies for at-risk groups should not exceed 20 mg/L (ADWG). Sudden changes in sodium levels in raw water supplies should be advised to Queensland Health, as these can affect medical equipment.
Sulfate	<4 mg/L The concentration of sulfate in drinking water should not exceed 250 mg/L (ADWG, based on taste/aesthetic considerations). ADWG 2011 health guideline: <500mg/L
Dissolved oxygen	>80 % saturation
Pesticides	<0.1 µg/L for an individual compound <1.0 µg/L combined total for all compounds

Indicator	Water quality objective ¹
Nitrite	<0.2 mg/L
Nitrate	<1.0mg/L
Chloride	<40 mg/L
Phosphate	<0.3 mg/L
Iron	<0.8 mg/L
Manganese	<0.06 mg/L
Other indicators (including physico-chemical indicators)	Refer to ADWG

Notes:

1. All values are based on advice/historical data provided by Mackay Water, except where otherwise indicated.
2. Sudden changes in sodium levels impact on medical equipment use, operation and calibration. Queensland Health should be advised of any such changes. The ADWG notes that 50 mg/L is a 'typical value' in reticulated supplies. The ADWG value for sodium is 180 mg/L (based on level at which taste become appreciable) however 'sodium salts cannot be easily removed from drinking water' and 'water authorities are strongly encouraged to keep sodium concentrations as low as possible'. It further notes that 'medical practitioners treating people with severe hypertension or congestive heart failure should be aware if the sodium concentration in the patient's drinking water exceeds 20 mg/L' (ADWG; sodium factsheet). The US EPA (2012 Drinking Water Standards and Health Advisories) health based value for sodium is 20 mg/L (for individuals on a 500 mg/day restricted sodium diet).

Sources: Mackay Water, Whitsunday Regional Council, Qld Health, Queensland Water Supply Regulator, Australian Drinking Water Guidelines (NHMRC, 2011)

Table 5. Aquaculture EV: Water quality objectives for tropical aquaculture.

Water parameter	Recommended range		Water parameter	Recommended range
	Fresh water	Marine		General aquatic
Dissolved oxygen	>4 mg/L	>4 mg/L	Arsenic	<0.05 mg/L
Temperature °C	21–32	24–33	Cadmium	<0.003 mg/L
pH	6.8–9.5	7–9.0	Calcium/Magnesium	10–160 mg/L
Ammonia (TAN, total ammonia-nitrogen)	<1.0 mg/L	<1.0 mg/L	Chromium	<0.1 mg/L
Ammonia (NH ₃ , un-ionised form)	<0.1 mg/L	<0.1 mg/L	Copper	<0.006 mg/L in soft water
Nitrate (NO ₃)	1–100 mg/L	1–100 mg/L	Cyanide	<0.005 mg/L
Nitrite (NO ₂)	<0.1 mg/L	<1.0 mg/L	Iron	<0.5 mg/L
Salinity	0–5 ppt	15–35 ppt	Lead	<0.03 mg/L
Hardness	20–450 mg/L		Manganese	<0.01 mg/L
Alkalinity	20–400 mg/L	>100 mg/L	Mercury	<0.00005 mg/L
Turbidity	<80 NTU		Nickel	<0.01 mg/L in soft water <0.04 mg/L in hard water
Chlorine	<0.003 mg/L		Tin	<0.001 mg/L
Hydrogen sulphide	<0.002 mg/L		Zinc	0.03–0.06 mg/L in soft water 1–2 mg/L in hard water

Source: Department of Primary Industries and Fisheries: Water Quality in Aquaculture—DPI Notes April 2004.

Table 6. Aquaculture EV: Water quality objectives for optimal growth of particular species in fresh water.

Water parameter	Barramundi	Eel	Silver perch	Jade perch	Sleepy cod	Redclaw
Dissolved oxygen	4–9 mg/L	>3 mg/L	>4 mg/L	>3 mg/L	>4.0 mg/L	>4.0 mg/L
Temperature °C	26–32	23–28	23–28	23–28	22–31	23–31
pH	7.5–8.5	7.0–8.5	6.5–9	6.5–9	7.0–8.5	7.0–8.5
Ammonia (TAN, Total ammonia-nitrogen)		<1.0 mg/L			<1.0 mg/L	<1.0 mg/L
Ammonia (NH ₃ , un-ionised form)*pH dependent.	<0.46 mg/L	<0.1 mg/L	<0.1 mg/L	<0.1 mg/L	<0.1 mg/L	<0.1 mg/L
Nitrate (NO ₃)			<100 mg/L			
Nitrite (NO ₂)	<1.5 mg/L	<1.0 mg/L	<0.1 mg/L		<1.0 mg/L	<1.0 mg/L
Salinity (extended periods)	0–35 ppt		<5 ppt	<5 ppt		<4 ppt
Salinity bath	0–35 ppt		5–10 ppt for 1 hour		max. 20 ppt for one hour	
Hardness (CaCO ₃)			>50 mg/L	>50 mg/L	>40 mg/L	>40 mg/L
Alkalinity	>20 mg/L		100–400 ppm	100–400 ppm	>40 mg/L	>40 mg/L
Chlorine	<0.04 mg/L				<0.04 mg/L	
Hydrogen sulphide	0–0.3 mg/L				0–0.3 mg/L	
Iron	<0.1 mg/L		<0.5 mg/L	<0.5 mg/L	<0.1 mg/L	<0.1 mg/L
Spawning temperature °C	Marine		23–28	23–28	>24 for more than three days	

Source: Department of Primary Industries and Fisheries: Water Quality in Aquaculture—DPI Notes April 2004.

Table 7. Aquaculture EV: Water quality objectives for optimal growth of particular marine species.

Water parameter	Barramundi		Tiger prawn		Kuruma prawn
	Hatchery	Grow out	Hatchery	Grow out	Grow out
Dissolved oxygen	Saturation	>4 mg/L	>4 mg/L	>3.5 mg/L	>4 mg/L
Temperature °C	28–30 optimum 25–31 range	28–30 optimum		26–32	24
pH	~8	~8	~8	7.5–8.5	7.5–8.5
Ammonia (TAN, total ammonia-nitrogen)		0.1–0.5 mg/L			
Ammonia (NH ₃ , un-ionised form)	<0.1 mg/L	<0.1 mg/L	<0.1 mg/L	<0.1 mg/L	<0.1 mg/L
Nitrate (NO ₃)	<1.0 mg/L	<1.0 mg/L	<1.0 mg/L	<1.0 mg/L	<1.0 mg/L
Nitrite (NO ₂)	<0.2 mg/L	<1.0 mg/L	<0.2 mg/L	<0.2 mg/L	<0.2 mg/L
Salinity	28–31 ppt	0–35 ppt		10–25 ppt optimum	30–35 ppt optimum
Alkalinity		105–125 mg/L CaCO ₃			
Clarity				30–40 cm Secchi disk	30–40 cm Secchi disk
Hydrogen sulphide		<0.3 mg/L			
Iron		<0.02 mg/L		<1.0 mg/L	
Spawning temperature °C		28–32		27–32	

Source: Department of Primary Industries and Fisheries—Water Quality in Aquaculture—DPI Notes April 2004 (as amended).

Table 8. Irrigation EV: Water quality objectives for thermotolerant (faecal) coliforms in irrigation waters used for food and non-food crops¹.

Intended use	Median values of thermotolerant coliforms (colony forming units—cfu) ²
Raw human food crops in direct contact with irrigation water (e.g. via sprays, irrigation of salad vegetables)	<10 cfu/100 mL
Raw human food crops not in direct contact with irrigation water (edible product separated from contact with water, e.g. by peel, use of trickle irrigation); or crops sold to consumers cooked or processed	<1000 cfu/100 mL
Pasture and fodder for dairy animals (without withholding period)	<100 cfu/100 mL
Pasture and fodder for dairy animals (with withholding period of five days)	<1000 cfu/100 mL
Pasture and fodder (for grazing animals except pigs and dairy animals, i.e. cattle, sheep and goats)	<1000 cfu/100 mL
Silviculture, turf, cotton, etc. (restricted public access)	<10 000 cfu/100 mL

Notes:

1. Adapted from ARMCANZ, ANZECC and NHMRC (1999).
2. Refer to AWQG, Volume 1, Section 4.2.3.3 for advice on testing protocols.

Source: AWQG, Volume 1, Section 4.2.3.3, Table 4.2.2.

Table 9. Irrigation EV: Water quality objectives for heavy metals and metalloids in agricultural irrigation water¹—long-term trigger value (LTV), short-term trigger value (STV) and soil cumulative contamination loading limit (CCL).

Element	Soil cumulative contaminant loading limit (CCL) ² (kg/ha)	Long-term trigger value (LTV) in irrigation water (up to 100 years) (mg/L)	Short-term trigger value (STV) in irrigation water (up to 20 years) (mg/L)
Aluminium	ND ²	5	20
Arsenic	20	0.1	2.0
Beryllium	ND	0.1	0.5
Boron	ND	0.5	Refer to AWQG, Vol 3, Table 9.2.18
Cadmium	2	0.01	0.05
Chromium	ND	0.1	1
Cobalt	ND	0.05	0.1
Copper	140	0.2	5
Fluoride	ND	1	2
Iron	ND	0.2	10
Lead	260	2	5
Lithium	ND	2.5 (0.075 for citrus crops)	2.5 (0.075 for citrus crops)
Manganese	ND	0.2	10
Mercury	2	0.002	0.002
Molybdenum	ND	0.01	0.05
Nickel	85	0.2	2
Selenium	10	0.02	0.05
Uranium	ND	0.01	0.1
Vanadium	ND	0.1	0.5
Zinc	300	2	5

Notes:

1. Concentrations in irrigation water should be less than the trigger values. Trigger values should only be used in conjunction with information on each individual element and the potential for off-site transport of contaminants (refer AWQG, Volume 3, Section 9.2.5).
2. ND = Not determined; insufficient background data to calculate CCL.

Source: AWQG, Volume 1, Section 4.2.6, Table 4.2.10.

Table 10. Stock watering EV: Water quality objectives for tolerances of livestock to total dissolved solids (salinity) in drinking water¹.

Livestock	Total dissolved solids (TDS) (mg/L)		
	No adverse effects on animals expected	Animals may have initial reluctance to drink or there may be some scouring, but stock should adapt without loss of production	Loss of production and decline in animal condition and health would be expected. Stock may tolerate these levels for short periods if introduced gradually
Beef cattle	0–4000	4000–5000	5000–10 000
Dairy cattle	0–2500	2500–4000	4000–7000
Sheep	0–5000	5000–10 000	10 000–13 000 ²
Horses	0–4000	4000–6000	6000–7000
Pigs	0–4000	4000–6000	6000–8000
Poultry	0–2000	2000–3000	3000–4000

Notes:

1. From ANZECC (1992), adapted to incorporate more recent information.
2. Sheep on lush green feed may tolerate up to 13 000 mg/L TDS without loss of condition or production.

Source: AWQG, Volume 1, Section 4.3.3.5, Table 4.3.1.

Table 11. Stock watering EV: Water quality objectives (low risk trigger values) for heavy metals and metalloids in livestock drinking water.

Metal or metalloid	Trigger value (low risk) ^{1,2} (mg/L)
Aluminium	5
Arsenic	0.5 (up to 5 ³)
Beryllium	ND
Boron	5
Cadmium	0.01
Chromium	1
Cobalt	1
Copper	0.4 (sheep), 1 (cattle), 5 (pigs), 5 (poultry)
Fluoride	2
Iron	not sufficiently toxic
Lead	0.1
Manganese	not sufficiently toxic
Mercury	0.002
Molybdenum	0.15
Nickel	1
Selenium	0.02
Uranium	0.2
Vanadium	ND
Zinc	20

Notes:

1. Higher concentrations may be tolerated in some situations (further details provided in AWQG, Volume 3, Section 9.3.5).
2. ND = not determined, insufficient background data to calculate.
3. May be tolerated if not provided as a food additive and natural levels in the diet are low.

Source: AWQG, Volume 1, Section 4.3.4, Table 4.3.2.

Table 12. Recreational waters: Alert levels and corresponding actions for management of cyanobacteria.

When cyanobacteria are present in large numbers they can present a significant hazard, particularly to primary contact users of waters. Water quality objectives for cyanobacteria in recreational waters are provided in Table 3. Monitoring/action requirements relative to cyanobacteria 'alert' levels are summarised below, and are explained more fully in the Guidelines for Managing Risks in Recreational Water (NHMRC, 2008). Further details on the process to determine suitability of waters for recreation, relative to historical cyanobacterial levels and susceptibility to cyanobacterial contamination, are contained in sections 6 and 7 of the NHMRC guidelines.

Green level surveillance mode ¹	Amber level alert mode ¹	Red level action mode ¹
Fresh waters		
≥ 500 to <5000 cells/mL <i>M. aeruginosa</i> or biovolume equivalent of >0.04 to <0.4 mm ³ /L for the combined total of all cyanobacteria.	≥ 5000 to <50 000 cells/mL <i>M. aeruginosa</i> or biovolume equivalent of ≥ 0.4 to <4 mm ³ /L for the combined total of all cyanobacteria where a known toxin producer is dominant in the total biovolume ² . or ³ ≥ 0.4 to <10 mm ³ /L for the combined total of all cyanobacteria where known toxin producers are not present.	Level 1 guideline ⁴ : ≥ 10 µg/L total microcystins or ≥ 50 000 cells/mL toxic <i>M. aeruginosa</i> or biovolume equivalent of ≥ 4 mm ³ /L for the combined total of all cyanobacteria where a known toxin producer is dominant in the total biovolume. or ³ Level 2 guideline ⁴ : ≥ 10 mm ³ /L for total biovolume of all cyanobacterial material where known toxins are not present. or cyanobacterial scums are consistently present ⁵ .
Coastal and estuarine waters		
<i>Karenia brevis</i>		
≤ 1 cell/mL	> 1– < 10 cells/mL	≥ 10 cells/mL
<i>Lyngbya majuscula</i> , <i>Pfiesteria</i> spp.		
History but no current presence of organism	Present in low numbers	Present in high numbers. (For <i>Lyngbya majuscula</i> this involves the relatively widespread visible presence of dislodged algal filaments in the water and washed up onto the beach)
<i>Nodularia spumigena</i> : See NHMRC, Chapter 6 (Cyanobacteria and algae in fresh water) for details.		

Notes:

- Recommended actions at different alert levels are outlined below (based on NHMRC, 2008, Table 6.6—fresh waters. Similar actions are outlined for coastal/estuarine waters in NHMRC Table 7.6):
 - Green:** Regular monitoring. Weekly sampling and cell counts at representative locations in the water body where known toxigenic species are present (i.e. *Microcystis aeruginosa*, *Anabaena circinalis*, *Cylindrospermopsis raciborskii*, *Aphanizomenon ovalisporum*, *Nodularia spumigena*); or fortnightly for other types including regular visual inspection of water surface for scums.
 - Amber:** Notify agencies as appropriate. Increase sampling frequency to twice weekly at representative locations in the water body where toxigenic species (above) are dominant within the alert level definition (i.e. total biovolume) to establish population growth and spatial variability in the water body. Monitor weekly or fortnightly where other types are dominant. Make regular visual inspections of water surface for scums. Decide on requirement for toxicity assessment or toxin monitoring.
 - Red:** Continue monitoring as for (amber) alert mode. Immediately notify health authorities for advice on health risk. ('In action mode the local authority and health authorities warn the public of the existence of potential health risks; for example, through the media and the erection of signs by the local authority.' NHMRC, 2008; 114). Make toxicity assessment or toxin measurement of water if this has not already been done. Health authorities warn of risk to public health (i.e. the authorities make a health risk assessment considering toxin monitoring data, sample type and variability).
- The definition of 'dominant' is where the known toxin producer comprises 75 per cent or more of the total biovolume of cyanobacteria in a representative sample.
- This applies where high cell densities or scums of 'non toxic' cyanobacteria are present i.e. where the cyanobacterial population has been tested and shown not to contain known toxins (microcystins, nodularian, cylindrospermopsin or saxitoxin).
- Health risks and levels: Level 1 is developed to protect against short-term health effects of exposure to cyanobacterial toxins ingested during recreational activity, whereas the Level 2 applies to the circumstance where there is a probability of increased likelihood of non-specific adverse health outcomes, principally respiratory, irritation and allergy symptoms, from exposure to very high cell densities of

cyanobacterial material irrespective of the presence of toxicity or known toxins (NHMRC, 2008;114).

5. This refers to the situation where scums occur at the recreation site each day when conditions are calm, particularly in the morning. Note that it is not likely that scums are always present and visible when there is a high population as the cells may mix down with wind and turbulence and then reform later when conditions become stable.

Source: Based on NHMRC (2008) Guideline for Managing Risks in Recreational Water (Tables 6.2, 6.6, 7.3).

4 Ways to improve water quality

The following documents are relevant in considering ways to improve water quality in the Proserpine River, Whitsunday Island and O'Connell River basins. The document list below is additional to the plans, guidelines and other sources referred to in previous sections, **and is provided for information only**.

Local plans, studies

- Council planning scheme and supporting codes, policies, available from relevant council websites.

Regional plans, studies

- Mackay, Isaac and Whitsunday Regional Plan, February 2012, available from the Department of State Development, Infrastructure and Planning.
- Drewry, J, Higham, W, Mitchell, C 2008, Water quality improvement plan (WQIP): Final report for the Mackay Whitsunday region, Mackay Whitsunday Natural Resource Management Group (Reef catchments).
- Whitsundays Plan of Management 2008, GBRMPA, available from the GBRMPA website.

State plans, policies, guidelines, agreements etc

- State Planning Policy 4/10: Healthy Waters, available from the department's website.
- Urban Stormwater Quality Planning Guidelines (Queensland Government), available from the department's website.
- Queensland Water Quality Guidelines (QWQG), available from the department's website.
- Water Resource (Whitsunday) Plan 2010, available from the Office of Queensland Parliamentary Counsel website, and supporting documents available from the Department of Natural Resources and Mines website.
- Monitoring and Sampling Manual, available from the department's website.
- Reef Water Quality Protection Plan, Australian and Queensland Governments, available from the Reef Water Quality Protection Plan's website.

Other supporting technical information – riparian management

- Managing riparian widths to achieve multiple objectives, fact sheet 13, Land and Water Australia, Australian Government, 2004.
- Improving water quality, fact sheet 3, Land & Water Australia, Australian Government, 2002.
- Riparian Land Management Technical Guidelines—Volume 1 and 2, November 1999, Land and Water Resources Research and Development Corporation (LWRRDC).
- Guidelines for Queensland Streambank Stabilisation with Riparian Vegetation, CRC for Catchment Hydrology, September 1999.
- Restoration of Fish Habitats—Fisheries Guidelines for Marine Areas, FHG002, available from the Department of Agriculture, Fisheries and Forestry.
- Fisheries Guidelines for Fish Habitat Buffer Zones, FHG003, available from the Department of Agriculture, Fisheries and Forestry.
- Guidelines for Riparian Filter Strips for Queensland Irrigators, CSIRO Land and Water, September 1999.

5 Dictionary

AMTD means the adopted middle thread distance which is the distance in kilometres, measured along the middle of a watercourse, that a specific point in the watercourse is from the watercourse's mouth or junction with the main watercourse (definition based on Water Regulation 2002).

ANZECC means the Australian and New Zealand Environment and Conservation Council.

Aquatic ecosystems (defined in the AWQG) comprise the animals, plants and micro-organisms that live in water, and the physical and chemical environment and climatic regime in which they interact. It is predominantly the physical components (e.g. light, temperature, mixing, flow, habitat) and chemical components (e.g. organic and inorganic carbon, oxygen, nutrients) of an ecosystem that determine what lives and breeds in it, and therefore the structure of the food web. Biological interactions (e.g. grazing and predation) can also play a part in structuring many aquatic ecosystems.

ARMCANZ means the Agriculture and Resource Management Council of Australia and New Zealand.

Basin means the basin name and number provided by Geoscience Australia, Canberra (3rd edition, 2004).

Biological integrity, of water, means the water's ability to support and maintain a balanced, integrative, adaptive community of organisms having a species composition, diversity and functional organisation comparable to that of the natural habitat of the locality in which the water is situated.

Biotoxin (defined in the AWQG): means a toxin (poison) which originates from a living thing (a plant, animal, fungi, bacteria, etc).

Catchment means the total area draining into a river, creek, reservoir or other body of water. The limits of a given catchment are the heights of land (such as hills or mountains) separating it from neighbouring catchments. Catchments can be made up of smaller subcatchments.

Ecological health (defined in the AWQG) means the 'health' or 'condition' of an ecosystem. It is the ability of an ecosystem to support and maintain key ecological processes and organisms so that their species compositions, diversity and functional organisations are as comparable as possible to those occurring in natural habitats within a region (also termed ecological integrity).

Environmental value (EV) means:

- (a) a quality or physical characteristic of the environment that is conducive to ecological health or public amenity or safety; or
- (b) another quality of the environment identified and declared to be an environmental value under an Environmental Protection Policy or Regulation (e.g. water suitable for swimming in or drinking).

The EVs for water that can be identified for protection are outlined in Table 13.

Highest astronomical tide (HAT) (defined in Marine Parks (Declaration) Regulation 2006) means the highest level of the tides that can be predicted to occur under average meteorological conditions and under any combination of astronomical conditions.

High water mark (defined in *Coastal Protection and Management Act 1995*) means the ordinary high water mark at spring tides.

Mean high water spring refer high water mark.

Queensland waters (as defined in *Acts Interpretation Act 1954*): means all waters that are a) within the limits of the State; or b) coastal waters of the State.

Sub-basin means part of a basin.

Subcatchment means part of a catchment.

Toxicant (defined in the AWQG): means a chemical capable of producing an adverse response (effect) in a biological system at concentrations that might be encountered in the environment, seriously injuring structure or function or producing death. Examples include pesticides, heavy metals and biotoxins.

Table 13. Suite of environmental values that can be chosen for protection.

Environmental values and definitions	ICON (as shown on plans)
<p>Aquatic ecosystem 'A community of organisms living within or adjacent to water, including riparian or foreshore area.' (EPP (Water), schedule 2 - Dictionary) The intrinsic value of aquatic ecosystems, habitat and wildlife in waterways and riparian areas, for example, biodiversity, ecological interactions, plants, animals, key species (such as turtles, platypus, seagrass and dugongs) and their habitat, food and drinking water. Waterways include perennial and intermittent surface waters, groundwaters, tidal and non-tidal waters, lakes, storages, reservoirs, dams, wetlands, swamps, marshes, lagoons, canals, natural and artificial channels and the bed and banks of waterways. (This EV incorporates the 'wildlife habitat' EV used in the South East Queensland Regional Water Quality Management Strategy). See below for more details on aquatic ecosystems, based on the EPP (Water).</p>	
<p>High ecological/conservation value waters 'Waters in which the biological integrity of the water is effectively unmodified or highly valued.' (EPP (Water), schedule 2).</p>	None
<p>Slightly disturbed waters 'Waters that have the biological integrity of high ecological value waters with slightly modified physical or chemical indicators but effectively unmodified biological indicators.' (EPP (Water), schedule 2).</p>	None
<p>Moderately disturbed waters 'Waters in which the biological integrity of the water is adversely affected by human activity to a relatively small but measurable degree.' (EPP (Water), schedule 2).</p>	None
<p>Highly disturbed waters 'Waters that are significantly degraded by human activity and have lower ecological value than high ecological value waters or slightly or moderately disturbed waters.' (EPP (Water), schedule 2).</p>	None
<p>Seagrass (goal within the aquatic ecosystem EV). Maintenance or rehabilitation of seagrass habitat. (Applies only to tidal waterways.)</p>	

67

Environmental values and definitions	ICON (as shown on plans)
<p>Irrigation Suitability of water supply for irrigation, for example, irrigation of crops, pastures, parks, gardens and recreational areas.</p>	
<p>Farm water supply/use Suitability of domestic farm water supply, other than drinking water. For example, water used for laundry and produce preparation.</p>	
<p>Stock watering Suitability of water supply for production of healthy livestock.</p>	
<p>Aquaculture Health of aquaculture species and humans consuming aquatic foods (such as fish, molluscs and crustaceans) from commercial ventures.</p>	
<p>Human consumers of aquatic foods Health of humans consuming aquatic foods, such as fish, crustaceans and shellfish from natural waterways. Note that in some areas oystering is a more specific goal identified under the human consumer EV (see below).</p>	
<p>Oystering (goal within the EV of human consumers of aquatic foods) Health of humans consuming oysters from natural waterways and commercial ventures. (Applies only to tidal waterways.)</p>	
<p>Primary recreation Health of humans during recreation which involves direct contact and a high probability of water being swallowed, for example, swimming, surfing, windsurfing, diving and water-skiing. Primary recreational use, of water, means full body contact with the water, including, for example, diving, swimming, surfing, waterskiing and windsurfing. (EPP (Water), s. 6).</p>	
<p>Secondary recreation Health of humans during recreation which involves indirect contact and a low probability of water being swallowed, for example, wading, boating, rowing and fishing. Secondary recreational use, of water, means contact other than full body contact with the water, including, for example, boating and fishing. (EPP (Water), s. 6).</p>	
<p>Visual recreation Amenity of waterways for recreation which does not involve any contact with water—for example, walking and picnicking adjacent to a waterway. Visual recreational use, of a water, means viewing the water without contact with it. (EPP (Water), s. 6).</p>	
<p>Drinking water supply Suitability of raw drinking water supply. This assumes minimal treatment of water is required, for example, coarse screening and/or disinfection.</p>	

Environmental values and definitions	ICON (as shown on plans)
<p>Industrial use Suitability of water supply for industrial use, for example, food, beverage, paper, petroleum and power industries, mining and minerals refining/processing. Industries usually treat water supplies to meet their needs.</p>	
<p>Cultural and spiritual values Indigenous and non-indigenous cultural heritage, for example:</p> <ul style="list-style-type: none"> • custodial, spiritual, cultural and traditional heritage, hunting, gathering and ritual responsibilities • symbols, landmarks and icons (such as waterways, turtles and frogs) • lifestyles (such as agriculture and fishing). <p>Cultural and spiritual values, of water, means its aesthetic, historical, scientific, social or other significance, to the present generation or past or future generations. (EPP (Water), s. 6).</p>	